

Systematic search for singularities in 3D Euler flows

Xinyu Zhao, Bartosz Protas

McMaster University

zhaox171@mcmaster.ca

<https://sites.google.com/view/zhaoxinyu>

3D Euler equations on a periodic domain

$$\begin{aligned}\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} &= -\nabla p, & (\mathbf{x}, t) \in \mathbb{T}^3 \times (0, T], \\ \nabla \cdot \mathbf{u} &= 0, & (\mathbf{x}, t) \in \mathbb{T}^3 \times (0, T], \\ \mathbf{u}|_{t=0} &= \boldsymbol{\eta}, & \mathbf{x} \in \mathbb{T}^3.\end{aligned}\tag{1}$$

- $\mathbf{u} = (u_1, u_2, u_3)$ is the velocity field.
- $p = p(\mathbf{x}, t)$ is the scalar pressure.
- $\mathbb{T}^3 = \mathbb{R}^3 / \mathbb{Z}^3$ is a unit cube.

3D Euler equations on a periodic domain

$$\begin{aligned}\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} &= -\nabla p, & (\mathbf{x}, t) \in \mathbb{T}^3 \times (0, T], \\ \nabla \cdot \mathbf{u} &= 0, & (\mathbf{x}, t) \in \mathbb{T}^3 \times (0, T], \\ \mathbf{u}|_{t=0} &= \boldsymbol{\eta}, & \mathbf{x} \in \mathbb{T}^3.\end{aligned}\tag{1}$$

- Local well-posedness has been established.

Theorem (Kato, 1972)

If $\boldsymbol{\eta} \in H^m(\mathbb{T}^3)$ or $H^m(\mathbb{R}^3)$ for some $m > 5/2$ and satisfies $\nabla \cdot \boldsymbol{\eta} = 0$, then there exists a time $T > 0$ such that (1) has a unique solution \mathbf{u} with the initial condition $\boldsymbol{\eta}$ and the solution satisfies $\mathbf{u} \in C([0, T]; H^m) \cap C^1([0, T]; H^{m-1})$.

3D Euler equations on a periodic domain

$$\begin{aligned}\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} &= -\nabla p, & (\mathbf{x}, t) &\in \mathbb{T}^3 \times (0, T], \\ \nabla \cdot \mathbf{u} &= 0, & (\mathbf{x}, t) &\in \mathbb{T}^3 \times (0, T], \\ \mathbf{u}|_{t=0} &= \boldsymbol{\eta}, & \mathbf{x} &\in \mathbb{T}^3.\end{aligned}\tag{1}$$

Open question (global well-posedness)

Does there exist a smooth initial condition $\boldsymbol{\eta} \in H^m$ for $m > 5/2$, such that

$$\lim_{t \rightarrow T^*} \|\mathbf{u}(t; \boldsymbol{\eta})\|_{H^m} = \infty, \quad 0 < T^* < \infty?$$

- The global-wellposedness of solutions of the Navier-Stokes equations is one of the “millennium problems” posed by the Clay Mathematics Institute.

3D Euler equations on a periodic domain

$$\begin{aligned}\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} &= -\nabla p, & (\mathbf{x}, t) &\in \mathbb{T}^3 \times (0, T], \\ \nabla \cdot \mathbf{u} &= 0, & (\mathbf{x}, t) &\in \mathbb{T}^3 \times (0, T], \\ \mathbf{u}|_{t=0} &= \boldsymbol{\eta}, & \mathbf{x} &\in \mathbb{T}^3.\end{aligned}\tag{1}$$

Beale-Kato-Majda (BKM) criterion (Beale et al. 1984)

A smooth solution \mathbf{u} of (1) develops a singularity at $t = T^*$ if and only if

$$\int_0^{T^*} \|\boldsymbol{\omega}(t)\|_{L^\infty} dt = \infty, \quad \boldsymbol{\omega} = \nabla \times \mathbf{u}.$$

- There have been various refinements of the BKM criterion (Kozono & Taniuchi 2000, Chae 2001).

3D Euler equations on a periodic domain

- Kerr 1993, Bustamante & Brachet 2012, Hou 2022, Kang et al. 2020, Kang & Protas 2022, etc.




Figure: (Hou & Li 2006) The 3D view of the vortex tube for $t = 0$ and $t = 6$. The tube is the isosurface at 60% of the maximum vorticity. The ribbons on the symmetry plane are the contours at other different values.

3D Euler equations on a periodic domain

$$\begin{aligned}\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} &= -\nabla p, & (\mathbf{x}, t) \in \mathbb{T}^3 \times (0, T], \\ \nabla \cdot \mathbf{u} &= 0, & (\mathbf{x}, t) \in \mathbb{T}^3 \times (0, T], \\ \mathbf{u}|_{t=0} &= \boldsymbol{\eta}, & \mathbf{x} \in \mathbb{T}^3.\end{aligned}\tag{1}$$

Open question (global well-posedness)

Does there exist a smooth initial condition $\boldsymbol{\eta} \in H^m$ for $m > 5/2$, such that

$$\lim_{t \rightarrow T^*} \|\mathbf{u}(t; \boldsymbol{\eta})\|_{H^m} = \infty, \quad 0 < T^* < \infty?$$

- We choose $m = 3$ and assume $\int_{\mathbb{T}^3} \mathbf{u} \, d\mathbf{x} = \mathbf{0}$, thus we consider the \dot{H}^3 -norm.

Optimization problem

- Objective functional

$$\Phi_T(\boldsymbol{\eta}) := \|\mathbf{u}(T; \boldsymbol{\eta})\|_{\dot{H}^3}^2, \quad \boldsymbol{\eta} \in \mathcal{M}_1.$$

$$\mathcal{M}_1 := \left\{ \boldsymbol{\eta} \in G^\sigma \mid \int_{\mathbb{T}^3} \boldsymbol{\eta} \, d\mathbf{x} = \mathbf{0}, \nabla \cdot \boldsymbol{\eta} = 0, \|\boldsymbol{\eta}\|_{\dot{H}^3} = 1 \right\}.$$

- G^σ (Gevrey space) is endowed with the inner product (Kukavica & Vicol 2009)

$$\langle \mathbf{v}, \mathbf{u} \rangle_{G^\sigma} = \sum_{\mathbf{j} \in \mathbb{Z}^3} (1 + |2\pi \mathbf{j}|^2)^3 e^{4\pi\sigma|\mathbf{j}|} \hat{\mathbf{v}}_{\mathbf{j}} \cdot \overline{\hat{\mathbf{u}}_{\mathbf{j}}}.$$

- Symmetry of the Euler equations: $\lambda \mathbf{u}(\mathbf{x}, \lambda t) \Rightarrow \|\boldsymbol{\eta}\|_{\dot{H}^3} = 1$.

Optimization problem

- Objective functional

$$\Phi_T(\boldsymbol{\eta}) := \|\mathbf{u}(T; \boldsymbol{\eta})\|_{\dot{H}^3}^2, \quad \boldsymbol{\eta} \in \mathcal{M}_1.$$

$$\mathcal{M}_1 := \left\{ \boldsymbol{\eta} \in G^\sigma \left| \int_{\mathbb{T}^3} \boldsymbol{\eta} \, d\mathbf{x} = \mathbf{0}, \nabla \cdot \boldsymbol{\eta} = 0, \|\boldsymbol{\eta}\|_{\dot{H}^3} = 1 \right. \right\}.$$

- Optimization problem: given $T \in \mathbb{R}_+$, find

$$\tilde{\boldsymbol{\eta}}_T = \operatorname{argmax}_{\boldsymbol{\eta} \in \mathcal{M}_1} \Phi_T(\boldsymbol{\eta}).$$

Evaluation of the gradient

- Objective functional: $\Phi_T(\boldsymbol{\eta}) := \|\mathbf{u}(T; \boldsymbol{\eta})\|_{\dot{H}^3}^2$.

$$\begin{aligned}\Phi'_T(\boldsymbol{\eta}; \boldsymbol{\eta}') &:= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\Phi_T(\boldsymbol{\eta} + \epsilon \boldsymbol{\eta}') - \Phi_T(\boldsymbol{\eta})] = \langle \nabla \Phi_T(\boldsymbol{\eta}), \boldsymbol{\eta}' \rangle_{G^\sigma} \\ &= 2 \langle \mathbf{u}(\cdot, T), \mathbf{u}'(\cdot, T) \rangle_{\dot{H}^3}.\end{aligned}$$

- Linearization of the Euler equations around the solution $(\mathbf{u}(t; \boldsymbol{\eta}), p)$

$$\begin{aligned}\mathcal{L} \begin{bmatrix} \mathbf{u}' \\ p' \end{bmatrix} &:= \begin{bmatrix} \partial_t \mathbf{u}' + \mathbf{u}' \cdot \nabla \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u}' + \nabla p' \\ \nabla \cdot \mathbf{u}' \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \\ \mathbf{u}'(0) &= \boldsymbol{\eta}'.\end{aligned}$$

Evaluation of the gradient

- Objective functional: $\Phi_T(\boldsymbol{\eta}) := \|\mathbf{u}(T; \boldsymbol{\eta})\|_{\dot{H}^3}^2$.

$$\begin{aligned}\Phi'_T(\boldsymbol{\eta}; \boldsymbol{\eta}') &:= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\Phi_T(\boldsymbol{\eta} + \epsilon \boldsymbol{\eta}') - \Phi_T(\boldsymbol{\eta})] = \langle \nabla \Phi_T(\boldsymbol{\eta}), \boldsymbol{\eta}' \rangle_{G^\sigma} \\ &= 2 \langle \mathbf{u}(\cdot, T), \mathbf{u}'(\cdot, T) \rangle_{\dot{H}^3} .\end{aligned}$$

- Adjoint states (\mathbf{u}^*, p^*)

$$\begin{aligned}\left(\mathcal{L} \begin{bmatrix} \mathbf{u}' \\ p' \end{bmatrix}, \begin{bmatrix} \mathbf{u}^* \\ p^* \end{bmatrix} \right) &:= \int_0^T \int_{\mathbb{T}^3} \mathcal{L} \begin{bmatrix} \mathbf{u}' \\ p' \end{bmatrix} \cdot \begin{bmatrix} \mathbf{u}^* \\ p^* \end{bmatrix} d\mathbf{x} dt \\ &= \left(\begin{bmatrix} \mathbf{u}' \\ p' \end{bmatrix}, \mathcal{L}^* \begin{bmatrix} \mathbf{u}^* \\ p^* \end{bmatrix} \right) + \int_{\mathbb{T}^3} \mathbf{u}^*(\mathbf{x}, T) \cdot \mathbf{u}'(\mathbf{x}, T) d\mathbf{x} \\ &\quad - \int_{\mathbb{T}^3} \mathbf{u}^*(\mathbf{x}, 0) \cdot \boldsymbol{\eta}'(\mathbf{x}) d\mathbf{x} \\ &= 0.\end{aligned}$$

Evaluation of the gradient

- Objective functional: $\Phi_T(\boldsymbol{\eta}) := \|\mathbf{u}(T; \boldsymbol{\eta})\|_{\dot{H}^3}^2$.

$$\begin{aligned}\Phi'_T(\boldsymbol{\eta}; \boldsymbol{\eta}') &= \langle \nabla \Phi_T(\boldsymbol{\eta}), \boldsymbol{\eta}' \rangle_{G^\sigma} \\ &= 2 \langle \mathbf{u}(\cdot, T), \mathbf{u}'(\cdot, T) \rangle_{\dot{H}^3} = 2 \langle |D|^6 \mathbf{u}(\cdot, T), \mathbf{u}'(\cdot, T) \rangle_{L^2} \\ &= \langle \mathbf{u}^*(0), \boldsymbol{\eta}' \rangle_{L^2},\end{aligned}$$

where $|D| = \sqrt{-\Delta}$.

- $\nabla \Phi_T(\boldsymbol{\eta}) = (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}^*(0)$.
- Adjoint system


$$\begin{aligned}\mathcal{L}^* \begin{bmatrix} \mathbf{u}^* \\ p^* \end{bmatrix} &:= \begin{bmatrix} -\partial_t \mathbf{u}^* - (\nabla \mathbf{u}^* + (\nabla \mathbf{u}^*)^T) \mathbf{u} - \nabla p^* \\ -\nabla \cdot \mathbf{u}^* \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \\ \mathbf{u}^*(T) &= 2|D|^6 \mathbf{u}(T).\end{aligned}$$

Riemannian conjugate gradient method

$$\boldsymbol{\eta}^{(n+1)} = \mathbf{R} \left[\boldsymbol{\eta}^{(n)} + \tau_n \mathbf{d}^{(n)} \right], \quad n = 0, 1, 2, \dots,$$

$$\boldsymbol{\eta}^{(0)} = \boldsymbol{\eta}_0. \quad \Leftarrow \text{Taylor-Green, Random, Hou 2022, Kerr 1993}$$

- $\mathbf{R}(\mathbf{v}) = \mathbf{v} / \|\mathbf{v}\|_{\dot{H}^3}, \quad \mathbf{v} \neq \mathbf{0}.$
- $\mathbf{d}^{(n)} = \mathbf{P}_{\mathcal{T}_n} \mathbf{G}^{(n)} + \beta_n \Gamma_{\tau_{n-1}} \mathbf{d}^{(n-1)} (\mathbf{d}^{(n-1)}), \quad n \geq 1, \quad \mathbf{G}^{(n)} := \nabla \Phi_T (\boldsymbol{\eta}^{(n)}),$
where β_n is computed using the Polak-Ribière approach.



Riemannian conjugate gradient method

$$\boldsymbol{\eta}^{(n+1)} = \mathbf{R} \left[\boldsymbol{\eta}^{(n)} + \tau_n \mathbf{d}^{(n)} \right], \quad n = 0, 1, 2, \dots,$$

$$\boldsymbol{\eta}^{(0)} = \boldsymbol{\eta}_0. \quad \Leftarrow \text{Taylor-Green, Random, Hou 2022, Kerr 1993}$$


- $\mathbf{R}(\mathbf{v}) = \mathbf{v} / \|\mathbf{v}\|_{\dot{H}^3}, \quad \mathbf{v} \neq \mathbf{0}.$
- $\mathbf{d}^{(n)} = \mathbf{P}_{\mathcal{T}_n} \mathbf{G}^{(n)} + \beta_n \Gamma_{\tau_{n-1}} \mathbf{d}^{(n-1)}, \quad n \geq 1, \quad \mathbf{G}^{(n)} := \nabla \Phi_T (\boldsymbol{\eta}^{(n)}),$
where β_n is computed using the Polak-Ribière approach.
- $\tau_n = \operatorname{argmax}_{\tau > 0} \left\{ \Phi_T \left(\mathbf{R} \left[\boldsymbol{\eta}^{(n)} + \tau \mathbf{d}^{(n)} \right] \right) \right\}.$

Numerical results


We use uniform meshes with N^3 grid points and denote the solution we obtain at time T by $\tilde{\boldsymbol{\eta}}_T^N$.

- Shorter times T : $\lim_{N \rightarrow \infty} \Phi_T \left(\tilde{\boldsymbol{\eta}}_T^N \right) < \infty$
 \implies equation is well-posed on $[0, T]$;
- Longer times T : $\lim_{N \rightarrow \infty} \Phi_T \left(\tilde{\boldsymbol{\eta}}_T^N \right) = \infty$
 \implies there is a possible singularity at $T^* \leq T$.
- In practice, we use $N^3 = 128^3, 256^3, 512^3, 1024^3$.

Different initial guesses




(a) $T = 25$




(b) $T = 75$

Figure: Dependence of the objective functional $\Phi_T(\eta^{(n)})$ on the iteration index n for different initial guesses.

Behavior of Φ_T for $T = 25$ and $T = 75$




(a) $T = 25$




(b) $T = 75$

Figure: Comparison of the objective functional at different spatial resolutions for different time windows.


Vorticity components of $\tilde{\eta}_{75}^{1024}$



(a)




(b)




(c)

Vorticity field of $u(75; \tilde{\eta}_{75}^{1024})$



(a)



(b)

Figure: Visualization of the vorticity field.

Time evolution of $|\omega(t; \tilde{\eta}_{75}^{1024})|$




Figure: Time evolution of the vorticity field.

Time evolution of $\omega^\perp(t; \tilde{\eta}_{75}^{1024})$




Figure: Time evolution of the vorticity component orthogonal to the symmetry plane.



Xinyu Zhao, Bartosz Protas

Systematic search for singularities in 3D Euler flows

J. Nonlinear Sci., 2023, 33(6)

Thank you!

Time evolution of $|\omega(t; \tilde{\eta}_{75}^{1024})|$






Figure: Time evolution of the vorticity field.

Numerical results for the long time window ($T = 75$)




(a)




(b)

Figure: Comparison of the time evolution of the \dot{H}^3 -norm and the maximum vorticity of solutions with different initial conditions.

Energy spectrum of $\mathbf{u}^N \left(t, \tilde{\boldsymbol{\eta}}_{25}^N \right)$ at $t = 0$ and $t = 25$




(a)




(b)

$$e(k,t) := \frac{1}{2} \sum_{j \leq |j| < j+1} |\hat{\mathbf{u}}_j(t)|^2, \quad k = 2\pi j, \quad j \in \mathbb{N}.$$

Energy spectrum of $\mathbf{u}^N \left(t, \tilde{\boldsymbol{\eta}}_{75}^N \right)$ at $t = 0$ and $t = 75$



(a)




(b)


$$e(k,t) = \frac{1}{2} \sum_{j \leq |j| < j+1} |\hat{\mathbf{u}}_j(t)|^2, \quad k = 2\pi j, \quad j \in \mathbb{N}.$$

Numerical results for the long time window ($T = 75$)

$$\frac{d\|\mathbf{u}(t)\|_{\dot{H}^3}}{dt} = C\|\mathbf{u}(t)\|_{\dot{H}^3}^{\alpha} \implies \ln\left(\frac{d\|\mathbf{u}(t)\|_{\dot{H}^3}}{dt}\right) = \ln(C) + \alpha \ln(\|\mathbf{u}(t)\|_{\dot{H}^3}).$$



(a)



(b)

Figure: Growth rate of $\|\mathbf{u}(t)\|_{\dot{H}^3}$.