

Rational points on curves and tropical geometry.

David Zureick-Brown (Emory University)
Eric Katz (Waterloo University)

Slides available at <http://www.mathcs.emory.edu/~dzb/slides/>

Specialization of Linear Series for Algebraic and Tropical Curves
BIRS

April 3, 2014

Faltings' theorem

Theorem (Faltings)

Let X be a smooth curve over \mathbb{Q} with genus at least 2. Then $X(\mathbb{Q})$ is finite.

Example

For $g \geq 2$, $y^2 = x^{2g+1} + 1$ has only finitely many solutions with $x, y \in \mathbb{Q}$.

Problem

- 1 Given X , compute $X(\mathbb{Q})$ exactly.
- 2 Compute bounds on $\#X(\mathbb{Q})$.

Conjecture (Uniformity)

There exists a constant $N(g)$ such that every smooth curve of genus g over \mathbb{Q} has at most $N(g)$ rational points.

Theorem (Caporaso, Harris, Mazur)

Lang's conjecture \Rightarrow *uniformity*.

Theorem (Coleman)

Let X be a curve of genus g and let $r = \text{rank}_{\mathbb{Z}} \text{Jac}_X(\mathbb{Q})$. Suppose $p > 2g$ is a prime of good reduction. Suppose $r < g$. Then

$$\#X(\mathbb{Q}) \leq \#X(\mathbb{F}_p) + 2g - 2.$$

Remark

- 1 A modified statement holds for $p \leq 2g$ or for $K \neq \mathbb{Q}$.
- 2 Note: **this does not prove uniformity** (since the first good p might be large).

Theorem (Stoll)

Let X be a curve of genus g and let $r = \text{rank}_{\mathbb{Z}} \text{Jac}_X(\mathbb{Q})$. Suppose $p > 2g$ is a prime of good reduction. Suppose $r < g$. Then

$$\#X(\mathbb{Q}) \leq \#X(\mathbb{F}_p) + 2r.$$

Bad reduction bound

Theorem (Lorenzini-Tucker, McCallum-Poonen)

Let X be a curve of genus g and let $r = \text{rank}_{\mathbb{Z}} \text{Jac}_X(\mathbb{Q})$. Suppose $p > 2g$ is a prime. Suppose $r < g$.

Let \mathcal{X} be a regular proper model of X . Then

$$\#X(\mathbb{Q}) \leq \#\mathcal{X}^{\text{sm}}(\mathbb{F}_p) + 2g - 2.$$

Remark

A recent improvement due to Stoll gives a uniform bound if $r \leq g - 3$ and X is hyperelliptic.

Theorem (Katz-ZB)

Let X be a curve of genus g and let $r = \text{rank}_{\mathbb{Z}} \text{Jac}_X(\mathbb{Q})$. Suppose $p > 2g$ is a prime. Let \mathcal{X} be a regular proper model of X . Suppose $r < g$. Then

$$\#X(\mathbb{Q}) \leq \#\mathcal{X}^{\text{sm}}(\mathbb{F}_p) + 2r.$$

Example (hyperelliptic curve with cuspidal reduction)

$$\begin{aligned} -2 \cdot 11 \cdot 19 \cdot 173 \cdot y^2 &= (x - 50)(x - 9)(x - 3)(x + 13)(x^3 + 2x^2 + 3x + 4) \\ &= x(x + 1)(x + 2)(x + 3)(x + 4)^3 \pmod{5}. \end{aligned}$$

Analysis

① $X(\mathbb{Q})$ contains

$$\{\infty, (50, 0), (9, 0), (3, 0), (-13, 0), (25, 20247920), (25, -20247920)\}$$

② $\#\mathcal{X}_5^{\text{sm}}(\mathbb{F}_5) = 5$

③ $7 \leq \#X(\mathbb{Q}) \leq \#\mathcal{X}_5^{\text{sm}}(\mathbb{F}_5) + 2 \cdot 1 = 7$


This determines $X(\mathbb{Q})$.

Non-example


$$\begin{aligned}y^2 &= x^6 + 5 \\ &= x^6 \pmod{5}.\end{aligned}$$

Analysis

- 1 $X(\mathbb{Q}) \supset \{\infty^+, \infty^-\}$
- 2 $\mathcal{X}^{\text{sm}}(\mathbb{F}_5) = \{\infty^+, \infty^-, \pm(1, \pm 1), \pm(2, \pm 2^3), \pm(3, \pm 3^3), \pm(4, \pm 4^3)\}$
- 3 $2 \leq \#X(\mathbb{Q}) \leq \#\mathcal{X}_5^{\text{sm}}(\mathbb{F}_5) + 2 \cdot 1 = 20$




$$\begin{aligned} y^2 &= x^6 + 5 \\ &= x^6 \pmod{5}. \end{aligned}$$



Note: no \mathbb{Z}_p -point can reduce to $(0, 0)$.

Models – not regular


$$\begin{aligned}y^2 &= x^6 + 5^2 \\ &= x^6 \pmod{5}\end{aligned}$$



Now: $(0, 5)$ reduces to $(0, 0)$.

Models – not regular (blow up)

$$\begin{aligned}y^2 &= x^6 + 5^2 \\ &= x^6 \pmod{5}\end{aligned}$$



Blow up.

Models – semistable example


$$\begin{aligned}y^2 &= (x(x-1)(x-2))^3 + 5 \\ &= x^6 \pmod{5}.\end{aligned}$$



Note: no point can reduce to $(0, 0)$. Local equation looks like $xy = 5$

Models – semistable example (not regular)


$$\begin{aligned}y^2 &= (x(x-1)(x-2))^3 + 5^4 \\ &= x^6 \pmod{5}\end{aligned}$$



Now: $(0, 5^2)$ reduces to $(0, 0)$. Local equation looks like $xy = 5^4$

Models – semistable example


$$\begin{aligned}y^2 &= (x(x-1)(x-2))^3 + 5^4 \\ &= x^6 \pmod{5}\end{aligned}$$



Blow up. Local equation looks like $xy = 5^3$

Models – semistable example (regular at $(0,0)$)

$$\begin{aligned}y^2 &= (x(x-1)(x-2))^3 + 5^4 \\ &= x^6 \pmod{5}\end{aligned}$$




Blow up. Local equation looks like $xy = 5$

Main Theorem

Theorem (Katz-ZB)

Let X be a curve of genus g and let $r = \text{rank}_{\mathbb{Z}} \text{Jac}_X(\mathbb{Q})$. Suppose $p > 2g$ is a prime. Let \mathcal{X} be a regular proper model of X . Suppose $r < g$. Then

$$\#X(\mathbb{Q}) \leq \#\mathcal{X}^{\text{sm}}(\mathbb{F}_p) + 2r.$$



(p -adic integration) There exists $V \subset H^0(X_{\mathbb{Q}_p}, \Omega_X^1)$ with $\dim_{\mathbb{Q}_p} V \geq g - r$ such that,

$$\int_P^Q \omega = 0 \quad \forall P, Q \in X(\mathbb{Q}), \omega \in V$$

(Coleman, via Newton Polygons) Number of zeroes in a residue disc D_P is $\leq 1 + n_P$, where $n_P = \#(\operatorname{div} \omega \cap D_P)$

(Riemann-Roch) $\sum n_P = 2g - 2$.

(Coleman's bound) $\sum_{P \in X(\mathbb{F}_p)} (1 + n_P) = \#X(\mathbb{F}_p) + 2g - 2$.

Example (from McCallum-Poonen's survey paper)

Example

$$X: y^2 = x^6 + 8x^5 + 22x^4 + 22x^3 + 5x^2 + 6x + 1$$

- ① Points reducing to $\tilde{Q} = (0, 1)$ are given by

$$x = p \cdot t, \text{ where } t \in \mathbb{Z}_p$$

$$y = \sqrt{x^6 + 8x^5 + 22x^4 + 22x^3 + 5x^2 + 6x + 1} = 1 + x^2 + \dots$$


②
$$\int_{(0,1)}^{P_t} \frac{xdx}{y} = \int_0^t (x - x^3 + \dots) dx$$

Stoll's idea: use multiple ω

(Coleman, via Newton Polygons) Number of zeroes of $\int \omega$ in a residue class D_P is $\leq 1 + n_P$, where $n_P = \#(\text{div } \omega \cap D_P)$

Let $\tilde{n}_P = \min_{\omega \in V} \#(\text{div } \omega \cap D_P)$

(2 examples) $r \leq g - 2$, $\omega_1, \omega_2 \in V$



(Stoll's bound) $\sum \tilde{n}_P \leq 2r$. (Recall $\dim_{\mathbb{Q}_p} V \geq g - r$)

Stoll's bound – proof ($D = \sum \tilde{n}_P P$)

(Wanted)

$$\dim H^0(X_{\mathbb{F}_p}, K - D) \geq g - r \Rightarrow \text{deg } D \leq 2r$$

(Clifford)

$$H^0(X_{\mathbb{F}_p}, K - D') \neq 0 \Rightarrow \dim H^0(X_{\mathbb{F}_p}, D') \leq \frac{1}{2} \text{deg } D' + 1$$

($D' = K - D$)

$$\dim H^0(X_{\mathbb{F}_p}, K - D) \leq \frac{1}{2} \text{deg}(K - D) + 1$$

(Assumption)

$$g - r \leq \dim H^0(X_{\mathbb{F}_p}, K - D)$$

(Recall $\dim_{\mathbb{Q}_p} V \geq g - r$)

Complications when $X_{\mathbb{F}_p}$ is singular

- 1 $\omega \in H^0(X, \Omega)$ may **vanish along components** of $X_{\mathbb{F}_p}$;
- 2 i.e. $H^0(X_{\mathbb{F}_p}, K - D) \neq 0 \not\Rightarrow D$ is special;
- 3 $\text{rank}(K - D) \neq \dim H^0(X_{\mathbb{F}_p}, K - D) - 1$

Summary

The relationship between $\dim H^0(X_{\mathbb{F}_p}, K - D)$ and $\deg D$ is less transparent and does not follow from geometric techniques.

Rank of a divisor

Definition (Rank of a divisor is)

- 1 $r(D) = -1$ if $|D|$ is empty.
- 2 $r(D) \geq 0$ if $|D|$ is nonempty
- 3 $r(D) \geq k$ if $|D - E|$ is nonempty for any effective E with $\deg E = k$.

Remark

- 1 If X is smooth, then $r(D) = \dim H^0(X, D) - 1$.
- 2 If X has multiple components, then $r(D) \neq \dim H^0(X, D) - 1$.

Remark

Ingredients of Stoll's proof only use formal properties of $r(D)$.

Need:


$$\text{(Clifford)} \quad r(K - D) \leq \frac{1}{2} \deg(K - D)$$

$$\text{(Large rank)} \quad r(K - D) \geq g - r - 1$$

$$\text{(Recall, } V \subset H^0(X_{\mathbb{Q}_p}, \Omega_X^1), \dim_{\mathbb{Q}_p} V \geq g - r)$$

Idea: any section $s \in H^0(X, D)$ can be scaled to not vanish on a component (but may now have zeroes or poles at other components.)



Divisors on graphs:



Semistable case

Idea: any section $s \in H^0(X, D)$ can be scaled to not vanish on a component (but may now have zeroes or poles at other components.)





Divisors on graphs:



Semistable case

Idea: any section $s \in H^0(X, D)$ can be scaled to not vanish on a component (but may now have zeroes or poles at other components.)

Divisors on graphs:




Divisors on graphs

Definition (Rank of a divisor is)

- 1 $r(D) = -1$ if $|D|$ is empty.
- 2 $r(D) \geq 0$ if $|D|$ is nonempty
- 3 $r(D) \geq k$ if $|D - E|$ is nonempty for any effective E with $\deg E = k$.

0




1

3

1

-1

1



1

0

Remark


$$r(D) \geq 0$$

Divisors on graphs

Definition (Rank of a divisor is)

- 1 $r(D) = -1$ if $|D|$ is empty.
- 2 $r(D) \geq 0$ if $|D|$ is nonempty
- 3 $r(D) \geq k$ if $|D - E|$ is nonempty for any effective E with $\deg E = k$.


0



3

1

1



1

-1

Remark

$$r(D) \geq 1$$

Semistable case – line bundles

Let \mathcal{X} be a curve over \mathbb{Z}_p with semistable special fiber $\mathcal{X}_{\mathbb{F}_p} = \bigcup X_i$.

Definition (Divisor associated to a line bundle)

Given $\mathcal{L} \in \text{Pic } \mathcal{X}$, define a divisor on Γ by

$$\sum_{v \in V(\Gamma)} (\deg \mathcal{L}_{X_i}) v_{X_i}.$$

Semistable case – line bundles



Let \mathcal{X} be a curve over \mathbb{Z}_p with semistable special fiber $\mathcal{X}_{\mathbb{F}_p} = \bigcup X_i$.

Definition (Divisor associated to a line bundle)

Given $\mathcal{L} \in \text{Pic } \mathcal{X}$, define a divisor on Γ by

$$\sum_{v \in V(\Gamma)} (\deg \mathcal{L}_{X_i}) v_{X_i}.$$

Example: $\mathcal{L} = \omega_{\mathcal{X}}$, $\mathcal{X}_{\mathbb{F}_p}$ totally degenerate ($g(X_i) = 0$)



Semistable case – line bundles



Let \mathcal{X} be a curve over \mathbb{Z}_p with semistable special fiber $\mathcal{X}_{\mathbb{F}_p} = \bigcup X_i$.

Definition (Divisor associated to a line bundle)

Given $\mathcal{L} \in \text{Pic } \mathcal{X}$, define a divisor on Γ by

$$\sum_{v \in V(\Gamma)} (\deg \mathcal{L}_{X_i}) v_{X_i}.$$

Example: $\mathcal{L} = \mathcal{O}(H)$ (H a “horizontal” divisor on \mathcal{X})



Semistable case – line bundles



Let \mathcal{X} be a curve over \mathbb{Z}_p with semistable special fiber $\mathcal{X}_{\mathbb{F}_p} = \bigcup X_i$.

Definition (Divisor associated to a line bundle)

Given $\mathcal{L} \in \text{Pic } \mathcal{X}$, define a divisor on Γ by

$$\sum_{v \in V(\Gamma)} (\deg \mathcal{L}_{X_i}) v_{X_i}.$$

Example: $\mathcal{L} = \mathcal{O}(X_i)$,



Definition

For $\bar{D} \in \text{Div } \Gamma$, $r_{\text{num}}(\bar{D}) \geq k$ if $|\bar{D} - \bar{E}|$ is non-empty for every effective \bar{E} of degree k .

Theorem (Baker, Norine)



Riemann-Roch for r_{num} .

Clifford's theorem for r_{num} .

Specialization: $r_{\text{num}}(\bar{D}) \geq r(D)$.

Formal corollary: $X(\mathbb{Q}) \leq \#\mathcal{X}^{\text{sm}}(\mathbb{F}_p) + 2r$ (for X totally degenerate).

Semistable case – main points



Remark (Main points)

- 1 Chip firing is the same as twisting by $\mathcal{O}(X_i)$.
- 2 If $\exists s \in H^0(\mathcal{X}, \mathcal{L})$ and $\text{div } s = \sum H_i + \sum n_i X_i$, then

$$\mathcal{L} \otimes \mathcal{O}(-n_1 X_1) \otimes \cdots \otimes \mathcal{O}(-n_k X_k)$$

specializes to an effective divisor on Γ .

- 3 The firing sequence (n_1, \dots, n_n) wins the chip firing game.

Semistable but not totally degenerate – abelian rank

Problems when $g(\Gamma) < g(X)$. (E.g. rank can increase after reduction.)



Definition (Abelian rank r_{ab})

Let $\mathcal{L} \in \mathcal{X}$ have specialization $D \in \text{Div } \Gamma$. Then $r_{ab}(\mathcal{L}) \geq k$ if

- 1 $|D - E|$ is nonempty for any effective E with $\deg E = k$, and
- 2 for every \mathcal{L}_E specializing to E , there exists some (n_1, \dots, n_k) such that

$$\mathcal{L}' := \mathcal{L} \otimes \mathcal{L}_E^{-1} \otimes \mathcal{O}(n_1 X_1) \otimes \cdots \otimes \mathcal{O}(n_k X_k)$$

has effective specialization and such that $H^0(X_i, \mathcal{L}'_{X_i}) \neq 0$ for every component X_i .



Theorem (Katz-ZB)

Clifford's theorem: $r_{\text{ab}}(K - D) \leq \frac{1}{2} \deg(K - D)$

Specialization: $r_{\text{ab}}(K - D) \geq g - r.$

Formal corollary: $X(\mathbb{Q}) \leq \#\mathcal{X}^{\text{sm}}(\mathbb{F}_p) + 2r$ (for semistable curves.)

Final remarks

Remark

Also prove: **semistable case** \Rightarrow **general case**.

Remark (Néron models)

- 1 Suppose $\mathcal{L} \in \text{Pic}_{\mathcal{X}}$ and $\deg(\mathcal{L}|_{\mathcal{X}_p}) = 0$.
- 2 $r_{\text{num}}(\mathcal{L}) = 0$ if and only if $\mathcal{L}|_{\mathcal{X}_p} \in \text{Pic}^0_{\mathcal{X}_p}$.
- 3 $r_{\text{ab}}(\mathcal{L}) = 0$ if and only if the image of $\mathcal{L}|_{\mathcal{X}_p}$ in $\text{Pic}^0_{\widetilde{\mathcal{X}_p}}$ is the identity.

Remark (Toric rank)

- 1 Can also define r_{tor} – additionally require that sections agree at nodes
- 2 r_{tor} incorporates the toric part of Néron model