

# The basic CSP reductions revisited

Libor Barto, joint with Michael Pinsker

Charles University in Prague

Banff workshop  
November 2014

## Outline

- ▶ Basic CSP reductions – 3 views
- ▶ Questions
- ▶ Basic CSP reductions revisited

## Notation

- ▶  $\mathcal{A}$  ... finite set of relations on  $A$
- ▶  $\mathbf{A}$  ... the clone of polymorphisms of  $\mathcal{A}$

## Basic reductions – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$

## Basic reductions – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$
- ▶  $\mathcal{B}$  is a “pp-power” of  $\mathcal{A}$   
ie.  $B = A^n$  and  
each  $l$ -ary  $R \in \mathcal{B}$  is pp-def from  $\mathcal{A}$  as an  $nl$ -ary relation

## Basic reductions – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$
- ▶  $\mathcal{B}$  is a “pp-power” of  $\mathcal{A}$   
ie.  $B = A^n$  and  
each  $l$ -ary  $R \in \mathcal{B}$  is pp-def from  $\mathcal{A}$  as an  $nl$ -ary relation
- ▶  $\mathcal{B} = \mathcal{A}|_S$  (induced substructure), where  $S$  is pp-def from  $\mathcal{A}$

## Basic reductions – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$
- ▶  $\mathcal{B}$  is a “pp-power” of  $\mathcal{A}$   
ie.  $B = A^n$  and  
each  $l$ -ary  $R \in \mathcal{B}$  is pp-def from  $\mathcal{A}$  as an  $nl$ -ary relation
- ▶  $\mathcal{B} = \mathcal{A}|_S$  (induced substructure), where  $S$  is pp-def from  $\mathcal{A}$
- ▶  $\mathcal{B} \cong \mathcal{A}/\sim$ , where  $\sim$  is pp-def from  $\mathcal{A}$

## Basic reductions – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$
- ▶  $\mathcal{B}$  is a “pp-power” of  $\mathcal{A}$   
ie.  $B = A^n$  and  
each  $l$ -ary  $R \in \mathcal{B}$  is pp-def from  $\mathcal{A}$  as an  $nl$ -ary relation
- ▶  $\mathcal{B} = \mathcal{A}|_S$  (induced substructure), where  $S$  is pp-def from  $\mathcal{A}$
- ▶  $\mathcal{B} \cong \mathcal{A}/\sim$ , where  $\sim$  is pp-def from  $\mathcal{A}$
- ▶ last three together  $\Leftrightarrow \mathcal{A}$  pp-interprets  $\mathcal{B}$

## Basic reductions – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$
- ▶  $\mathcal{B}$  is a “pp-power” of  $\mathcal{A}$   
ie.  $B = A^n$  and  
each  $l$ -ary  $R \in \mathcal{B}$  is pp-def from  $\mathcal{A}$  as an  $nl$ -ary relation
- ▶  $\mathcal{B} = \mathcal{A}|_S$  (induced substructure), where  $S$  is pp-def from  $\mathcal{A}$
- ▶  $\mathcal{B} \cong \mathcal{A}/\sim$ , where  $\sim$  is pp-def from  $\mathcal{A}$
- ▶ last three together  $\Leftrightarrow \mathcal{A}$  pp-interprets  $\mathcal{B}$
- ▶  $\mathcal{B}$  is homomorphically equivalent to  $\mathcal{A}$  ... WLOG core



## Basic reductions – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$
- ▶  $\mathcal{B}$  is a “pp-power” of  $\mathcal{A}$   
ie.  $B = A^n$  and  
each  $l$ -ary  $R \in \mathcal{B}$  is pp-def from  $\mathcal{A}$  as an  $nl$ -ary relation
- ▶  $\mathcal{B} = \mathcal{A}|_S$  (induced substructure), where  $S$  is pp-def from  $\mathcal{A}$
- ▶  $\mathcal{B} \cong \mathcal{A}/\sim$ , where  $\sim$  is pp-def from  $\mathcal{A}$
- ▶ last three together  $\Leftrightarrow \mathcal{A}$  pp-interprets  $\mathcal{B}$
  
- ▶  $\mathcal{B}$  is homomorphically equivalent to  $\mathcal{A}$  ... WLOG core
- ▶  $\mathcal{B} = \mathcal{A} \cup \{a\}$ , if  $\mathcal{A}$  is a core

## Basic reductions – via algebraic constructions

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathbf{B}$  is an expansion of  $\mathbf{A}$  (ie.  $\mathbf{A} \subseteq \mathbf{B}$ )
- ▶  $\mathbf{B} = \mathbf{A}^n$
- ▶  $\mathbf{B}$  is a subalgebra of  $\mathbf{A}$
- ▶  $\mathbf{B} \cong \mathbf{A}/\sim$ , where  $\sim$  is a congruence of  $\mathbf{A}$
- ▶ last three  $\Leftrightarrow \mathbf{B} \in \text{HSP}^{\text{fin}}(\mathbf{A})$
  
- ▶ Finite: WLOG  $\mathbf{A}$  idempotent
- ▶ Infinite: A bit different

## Definition

$\xi : \mathbf{A} \rightarrow \mathbf{B}$  is a **clone homomorphism**, if it

- ▶ preserves arities
- ▶ sends projections to projections  $\xi(\pi_i^n) = \pi_i^n$
- ▶ preserves composition:  
$$\xi(f(g_1, \dots, g_n)) = \xi(f)(\xi(g_1), \dots, \xi(g_n)),$$
where  $f \in \mathbf{A}$  is  $n$ -ary,  $g_i \in \mathbf{A}$  is  $m$ -ary

Alternatively:  $\xi$ -images satisfy the same identities.

# The 3 together for finite

Theorem (Bodnaruk et al./Geiger; Birkhoff)

TFAE if  $A, B$  are finite:

1.  $A$  pp-interprets  $B$

$$A \xrightarrow{\text{pp-power}} \mathcal{E} \xrightarrow{\text{substr}} \mathcal{F} \xrightarrow{\text{quotient}} B$$

2.  $B$  is an expansion of a clone in  $\text{HSP}^{\text{fin}}(\mathbf{A})$

$$\mathbf{A} \xrightarrow{\sim} \mathbf{A}^n \xrightarrow{\text{subalg}} \mathbf{C} \xrightarrow{\text{quotient}} \mathbf{C}/\sim \xrightarrow{\text{expansion}} \mathbf{B}$$

3. There exists a clone homomorphism  $\xi : \mathbf{A} \rightarrow \mathbf{B}$

### (3) $\Rightarrow$ (2)

Assume  $\xi : \mathbf{A} \rightarrow \mathbf{B}$  is a clone homomorphism.

Want:

$$\mathbf{A} \rightsquigarrow \mathbf{A}^n \xrightarrow{\text{subalg}} \mathbf{C} \xrightarrow{\text{quotient}} \mathbf{C}/\sim \xrightarrow{\text{expansion}} \mathbf{B}$$

- ▶ Say  $B = \{b_1, \dots, b_k\}$
- ▶  $n = A^k$
- ▶  $C = k$ -ary operations in  $\mathbf{A}$   
( $\mathbf{C}$  is the free algebra with  $B$  generators)
- ▶ Define  $f : C \rightarrow B$  by  $t \mapsto \xi(t)(b_1, \dots, b_k)$
- ▶  $\sim = \ker f$  is a congruence of  $\mathbf{A}$  and  $f$  gives an isomorphism  $\mathbf{C}/\sim \rightarrow \xi(\mathbf{A})$

# The 3 together for infinite

Theorem (Bodirsky, Nešetřil; Bodrisky, Pisker)

TFAE if  $\mathcal{A}, \mathcal{B}$   $\omega$ -categorical:

1.  $\mathcal{A}$  pp-interprets  $\mathcal{B}$

$$\mathcal{A} \xrightarrow[\text{pp-power}]{\rightsquigarrow} \mathcal{E} \xrightarrow[\text{substr}]{\rightsquigarrow} \mathcal{F} \xrightarrow[\text{quotient}]{\rightsquigarrow} \mathcal{B}$$

2.  $\mathcal{B}$  is an expansion of a clone in  $\text{HSP}^{\text{fin}}(\mathbf{A})$ .

$$\mathbf{A} \rightsquigarrow \mathbf{A}^n \xrightarrow[\text{subalg}]{\rightsquigarrow} \mathbf{C} \xrightarrow[\text{quotient}]{\rightsquigarrow} \mathbf{C}/\sim \xrightarrow[\text{expansion}]{\rightsquigarrow} \mathbf{B}$$

3. There exists a continuous clone homomorphism  $\xi : \mathbf{A} \rightarrow \mathbf{B}$  such that  $\overline{\xi(\mathbf{A})}$  is oligomorphic

# Algebraic dichotomy conjecture vs. reality

Conjecture (Bulatov, Jeavons, Krokhin; Barto, Kozik)

Assume  $A$  finite. TFAE

- ▶  $\text{CSP}(\mathcal{A})$  in P
- ▶  $\mathbf{A}$  contains an operation  $t$  of arity  $\geq 2$  such that

$$t(x_1, x_2, \dots, x_n) = t(x_2, \dots, x_n, x_1)$$

Reality can be worse:

## Future theorem (Antonín Barto, Bálint Maróti 2013)

Assume  $A$  finite. TFAE

- ▶  $\text{CSP}(A)$  in  $P$
- ▶  $A$  contains an operations  $t_1, t_2, \dots$  such that

$$t_1(x_1, t_2(x_3, x_2), t_3(x_{123})) = t_3(t_3(t_2(x_{13}, x_2)))$$

$$t_{20}(t_{12}(x_1), x_1, x_2) = t_{13}(x_2, x_1)$$

...

...but certainly the characterization looks like this (for any complexity class)



- ▶ Theorem linking the 3 views does not cover all the easy reductions...

# Questions

- ▶ Theorem linking the 3 views does not cover all the easy reductions...
- ▶ ...and by adding homomorphic equivalence we get an essentially coarser ordering. What is this ordering?

# Questions

- ▶ Theorem linking the 3 views does not cover all the easy reductions...
- ▶ ...and by adding homomorphic equivalence we get an essentially coarser ordering. What is this ordering?
- ▶ In most identities relevant in CSP, there are no nested terms. Is it possible to prove that nesting is not necessary?

## Basic reductions revisited – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$
- ▶  $\mathcal{B}$  is a pp-power of  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A}|_{\mathcal{S}}$
- ▶  $\mathcal{B} = \mathcal{A}/\sim$
- ▶  $\mathcal{B}$  is homomorphically equivalent to  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A} \cup \{a\}$ , if  $\mathcal{A}$  is a core

## Basic reductions revisited – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$  **redundant**
- ▶  $\mathcal{B}$  is a pp-power of  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A}|_{\mathcal{S}}$
- ▶  $\mathcal{B} = \mathcal{A}/\sim$
- ▶  $\mathcal{B}$  is homomorphically equivalent to  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A} \cup \{a\}$ , if  $\mathcal{A}$  is a core

## Basic reductions revisited – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$  **redundant**
- ▶  $\mathcal{B}$  is a pp-power of  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A}|_{\mathcal{S}}$  **redundant**
- ▶  $\mathcal{B} = \mathcal{A}/\sim$
- ▶  $\mathcal{B}$  is homomorphically equivalent to  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A} \cup \{a\}$ , if  $\mathcal{A}$  is a core

## Basic reductions revisited – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$  **redundant**
- ▶  $\mathcal{B}$  is a pp-power of  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A}|_{\mathcal{S}}$  **redundant**
- ▶  $\mathcal{B} = \mathcal{A}/\sim$  **redundant**
- ▶  $\mathcal{B}$  is homomorphically equivalent to  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A} \cup \{a\}$ , if  $\mathcal{A}$  is a core

## Basic reductions revisited – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$  **redundant**
- ▶  $\mathcal{B}$  is a pp-power of  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A}|_{\mathcal{S}}$  **redundant**
- ▶  $\mathcal{B} = \mathcal{A}/\sim$  **redundant**
- ▶  $\mathcal{B}$  is homomorphically equivalent to  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A} \cup \{a\}$ , if  $\mathcal{A}$  is a core **redundant** (less obvious)



# Basic reductions revisited – via relations

$\text{CSP}(\mathcal{B})$  is log-space reducible to  $\text{CSP}(\mathcal{A})$  if

- ▶  $\mathcal{B}$  is pp-definable from  $\mathcal{A}$  **redundant**
- ▶  $\mathcal{B}$  is a pp-power of  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A}|_{\mathcal{S}}$  **redundant**
- ▶  $\mathcal{B} = \mathcal{A}/\sim$  **redundant**
- ▶  $\mathcal{B}$  is homomorphically equivalent to  $\mathcal{A}$
- ▶  $\mathcal{B} = \mathcal{A} \cup \{a\}$ , if  $\mathcal{A}$  is a core **redundant** (less obvious)

## Observation

*TFAE*

- ▶  $\mathcal{B}$  can be obtained from  $\mathcal{A}$  using the above constructions
- ▶  $\mathcal{B}$  is homomorphically equivalent to a pp-power of  $\mathcal{A}$

## Basic reductions revisited – via algebraic constructions

- ▶  $\mathbf{B}$  is an expansion of  $\mathbf{A}$  (ie.  $\mathbf{A} \subseteq \mathbf{B}$ )
- ▶  $\mathbf{B} = \mathbf{A}^n$
- ▶  $\mathbf{B}$  is a subalgebra of  $\mathbf{A}$
- ▶  $\mathbf{B} = \mathbf{A} / \sim$ , where  $\sim$  is a congruence of  $\mathbf{A}$

# Basic reductions revisited – via algebraic constructions

- ▶  $\mathbf{B}$  is an expansion of  $\mathbf{A}$  (ie.  $\mathbf{A} \subseteq \mathbf{B}$ )
- ▶  $\mathbf{B} = \mathbf{A}^n$
- ▶  $\mathbf{B}$  is a subalgebra of  $\mathbf{A}$
- ▶  $\mathbf{B} = \mathbf{A} / \sim$ , where  $\sim$  is a congruence of  $\mathbf{A}$

## Definition

$\mathbf{B}$  is a **unary modification** of  $\mathbf{A}$  if  $\exists f : A \rightarrow B, \exists g : B \rightarrow A$ :  
 $\mathbf{B} = \langle \bar{t} : t \in \mathbf{A} \rangle, \quad \bar{t}(x_1, \dots, x_k) = f(t(g(x_1), \dots, g(x_k)))$

# Basic reductions revisited – via algebraic constructions

- ▶  $\mathbf{B}$  is an expansion of  $\mathbf{A}$  (ie.  $\mathbf{A} \subseteq \mathbf{B}$ )
- ▶  $\mathbf{B} = \mathbf{A}^n$
- ▶  $\mathbf{B}$  is a subalgebra of  $\mathbf{A}$  **redundant**
- ▶  $\mathbf{B} = \mathbf{A} / \sim$ , where  $\sim$  is a congruence of  $\mathbf{A}$

## Definition

$\mathbf{B}$  is a **unary modification** of  $\mathbf{A}$  if  $\exists f : A \rightarrow B, \exists g : B \rightarrow A$ :  
 $\mathbf{B} = \langle \bar{t} : t \in \mathbf{A} \rangle, \quad \bar{t}(x_1, \dots, x_k) = f(t(g(x_1), \dots, g(x_k)))$

# Basic reductions revisited – via algebraic constructions

- ▶  $\mathbf{B}$  is an expansion of  $\mathbf{A}$  (ie.  $\mathbf{A} \subseteq \mathbf{B}$ )
- ▶  $\mathbf{B} = \mathbf{A}^n$
- ▶  $\mathbf{B}$  is a subalgebra of  $\mathbf{A}$  **redundant**
- ▶  $\mathbf{B} = \mathbf{A} / \sim$ , where  $\sim$  is a congruence of  $\mathbf{A}$  **redundant**

## Definition

$\mathbf{B}$  is a **unary modification** of  $\mathbf{A}$  if  $\exists f : A \rightarrow B, \exists g : B \rightarrow A$ :  
 $\mathbf{B} = \langle \bar{t} : t \in \mathbf{A} \rangle, \quad \bar{t}(x_1, \dots, x_k) = f(t(g(x_1), \dots, g(x_k)))$

# Basic reductions revisited – via algebraic constructions

- ▶  $\mathbf{B}$  is an expansion of  $\mathbf{A}$  (ie.  $\mathbf{A} \subseteq \mathbf{B}$ )
- ▶  $\mathbf{B} = \mathbf{A}^n$
- ▶  $\mathbf{B}$  is a subalgebra of  $\mathbf{A}$  **redundant**
- ▶  $\mathbf{B} = \mathbf{A} / \sim$ , where  $\sim$  is a congruence of  $\mathbf{A}$  **redundant**

## Definition

$\mathbf{B}$  is a **unary modification** of  $\mathbf{A}$  if  $\exists f : A \rightarrow B, \exists g : B \rightarrow A$ :  
 $\mathbf{B} = \langle \bar{t} : t \in \mathbf{A} \rangle, \quad \bar{t}(x_1, \dots, x_k) = f(t(g(x_1), \dots, g(x_k)))$

## Observation

*TFAE for  $\omega$ -categorical*

- ▶  $\mathcal{B}$  is homomorphically equivalent to a pp-power of  $\mathcal{A}$
- ▶  $\mathbf{B}$  is an expansion of a unary modification of a power of  $\mathbf{A}$   
(finite: equivalently  $g$  can be taken injective)

## Definition

$\xi : \mathbf{A} \rightarrow \mathbf{B}$  is a **weak clone homomorphism**, if it

- ▶ preserves arities
- ▶ preserves composition with projections:

$$\xi(f(\pi_{I_1}, \dots, \pi_{I_n})) = \xi(f)(\pi_{I_1}, \dots, \pi_{I_n}), \text{ where } f \in \mathbf{A} \text{ is } n\text{-ary}$$

Alternatively:  $\xi$ -images satisfy the same strongly linear identities  
(=height 1 terms on both sides)

## Theorem

*TFAE if  $A, B$  are finite:*

1.  $B$  is homo equivalent to a pp-power of  $A$

$$A \overset{\text{pp-power}}{\rightsquigarrow} \mathcal{E} \overset{\text{homo-eq}}{\rightsquigarrow} B$$

2.  $B$  is expansion of a unary modification of a power of  $A$

$$A \rightsquigarrow A^n \overset{\text{unary mod}}{\rightsquigarrow} D \overset{\text{expansion}}{\rightsquigarrow} B$$

3. There exists a weak clone homomorphisms  $\xi : A \rightarrow B$



### (3) $\Rightarrow$ (2)

Assume  $\xi : \mathbf{A} \rightarrow \mathbf{B}$  is a clone homomorphism.

Want:

$$\mathbf{A} \rightsquigarrow \mathbf{A}^n \xrightarrow{\text{shrink}} \mathbf{D} \xrightarrow{\text{expansion}} \mathbf{B}$$

- ▶ Say  $B = \{b_1, \dots, b_k\}$
- ▶  $n = A^k$
- ▶  $C = k$ -ary operations in  $\mathbf{A}$
- ▶  $\mathbf{D} =$  the clone generated by  $\xi(\mathbf{A})$
- ▶ Define  $f : A^k \rightarrow B$  by  $t \mapsto \xi(t)(b_1, \dots, b_k)$  on  $C$ , otherwise arbitrary
- ▶ Define  $g : B \rightarrow A^k$  by  $b_i \mapsto \pi_i$
- ▶  $\mathbf{D}$  is the unary modification of  $\mathbf{A}^k$  given by  $f, g$  since  $\bar{t} = \xi(t)$

## Conjecture (Bulatov, Jeavons, Krokhin; Barto, Kozik)

Assume  $A$  finite. TFAE

- ▶  $\text{CSP}(\mathcal{A})$  in P
- ▶  $\mathbf{A}$  contains an operation  $t$  of arity  $\geq 2$  such that

$$t(x_1, x_2, \dots, x_n) = t(x_2, \dots, x_n, x_1)$$

Reality can be worse

Future theorem ([Anna Kozik, Hermenegilda Pinsker 2019](#))

*Assume  $A$  finite. TFAE*

- ▶  $\text{CSP}(A)$  in  $P$
- ▶  $A$  contains an operations  $t_1, t_2, \dots$  such that

$$t_1(x_3, x_1, x_2, x_2) = t_2(x_3, x_3, x_1)$$

$$t_{20}(x_1, x_1, x_2) = t_{13}(x_2, x_1)$$

...but certainly the characterization looks like this  
**(for any complexity class)**

# The 3 together revisited for infinite

## Theorem

*TFAE if  $\mathcal{A}, \mathcal{B}$   $\omega$ -categorical.*

- 1.  $\mathcal{B}$  is homo equivalent to a pp-power of  $\mathcal{A}$*
- 2.  $\mathbf{B}$  is expansion of a unary modification of a power of  $\mathbf{A}$*

*And these conditions are implied by*

- 3 There exists continuous  $\xi : \mathbf{A} \rightarrow \mathbf{B}$  which preserves arities and*

$$\xi(\alpha(t(\beta_1, \dots, \beta_n))) = \xi(\alpha)\xi(t)(\xi(\beta_1), \dots, \xi(\beta_n))$$

*where  $f \in \mathbf{A}$  is  $n$ -ary, and  $\alpha, \beta_i \in \mathbf{A}$  are unary bijections.*

**Remember:** The complexity of  $\text{CSP}(\mathcal{A})$  for finite  $A$  depends only on linear identities satisfied by  $\mathbf{A}$

**Remember:** The complexity of  $\text{CSP}(\mathcal{A})$  for finite  $A$  depends only on linear identities satisfied by  $\mathbf{A}$

**To do:**

- ▶ Can we use it?

**Remember:** The complexity of  $\text{CSP}(\mathcal{A})$  for finite  $A$  depends only on linear identities satisfied by  $\mathbf{A}$

**To do:**

- ▶ Can we use it?
- ▶ Improve the 3 views theorem for infinite domains

**Remember:** The complexity of  $\text{CSP}(\mathcal{A})$  for finite  $A$  depends only on linear identities satisfied by  $\mathbf{A}$

**To do:**

- ▶ Can we use it?
- ▶ Improve the 3 views theorem for infinite domains
- ▶ Basic reductions  $\rightarrow$  preorder such that  $\mathcal{A} \leq \mathcal{B}$  then  $\text{CSP}(\mathcal{A})$  is easier than  $\text{CSP}(\mathcal{B})$   
Can we find some more reductions?  
Optimally characterizing log-space reduction (optimistic)



**Remember:** The complexity of  $\text{CSP}(\mathcal{A})$  for finite  $A$  depends only on linear identities satisfied by  $\mathbf{A}$

**To do:**

- ▶ Can we use it?
- ▶ Improve the 3 views theorem for infinite domains
- ▶ Basic reductions  $\rightarrow$  preorder such that  $\mathcal{A} \leq \mathcal{B}$  then  $\text{CSP}(\mathcal{A})$  is easier than  $\text{CSP}(\mathcal{B})$   
Can we find some more reductions?  
Optimally characterizing log-space reduction (optimistic)

**Thank you!**