

ELLIPTIC DIMER MODELS
AND
GENUS 1 HARNACK CURVES

Béatrice de Tilière
University Paris-Dauphine

work in progress with

Cédric Boutillier (Sorbonne) & David Cimasoni (Genève)


Banff, November 21, 2019

OUTLINE

- Dimer model
- Dimer model and Harnack curves
- Minimal isoradial immersions
- Elliptic dimer model
- Results

DIMER MODEL: DEFINITION

- ▶ Planar, bipartite graph $G = (V = B \cup W, E)$.




- ▶ **Dimer configuration** M : subset of edges s.t. each vertex is incident to exactly one edge of $M \rightsquigarrow \mathcal{M}(G)$.
- ▶ **Positive weight function** on edges: $\nu = (\nu_e)_{e \in E}$.
- ▶ **Dimer Boltzmann measure** (G finite):

$$\forall M \in \mathcal{M}(G), \quad \mathbb{P}_{\text{dimer}}(M) = \frac{\prod_{e \in M} \nu_e}{Z_{\text{dimer}}(G, \nu)}.$$

where $Z_{\text{dimer}}(G, \nu)$ is the **dimer partition function**.

DIMER MODEL: DEFINITION

- ▶ Planar, bipartite graph $G = (V = B \cup W, E)$.



- ▶ **Dimer configuration** M : subset of edges s.t. each vertex is incident to exactly one edge of $M \rightsquigarrow \mathcal{M}(G)$.
- ▶ **Positive weight function** on edges: $\nu = (\nu_e)_{e \in E}$.
- ▶ **Dimer Boltzmann measure** (G finite):

$$\forall M \in \mathcal{M}(G), \quad \mathbb{P}_{\text{dimer}}(M) = \frac{\prod_{e \in M} \nu_e}{Z_{\text{dimer}}(G, \nu)}.$$

where $Z_{\text{dimer}}(G, \nu)$ is the **dimer partition function**.

DIMER MODEL: KASTELEYN MATRIX

► Kasteleyn matrix (Percus-Kuperberg version)

- Edge $wb \rightsquigarrow$ angle ϕ_{wb} s.t. for every face $w_1, b_1, \dots, w_k, b_k$:

$$\sum_{j=1}^k (\phi_{w_j b_j} - \phi_{w_{j+1} b_j}) \equiv (k-1)\pi \pmod{2\pi}.$$

- K is the corresponding **twisted adjacency matrix**.

$$K_{w,b} = \begin{cases} \nu_{wb} e^{i\phi_{wb}} & \text{if } w \sim b \\ 0 & \text{otherwise.} \end{cases}$$

DIMER MODEL: FOUNDING RESULTS

- ▶ Assume G finite.

THEOREM ([KASTELEYN'61] [KUPERBERG'98])

$$Z_{\text{dimer}}(G, \nu) = |\det(K)|.$$

THEOREM (KENYON'97)

Let $\mathcal{E} = \{e_1 = w_1 b_1, \dots, e_n = w_n b_n\}$ be a subset of edges of G , then:


$$\mathbb{P}_{\text{dimer}}(e_1, \dots, e_n) = \left| \left(\prod_{j=1}^n K_{w_j, b_j} \right) \det(K^{-1})_{\mathcal{E}} \right|,$$

where $(K^{-1})_{\mathcal{E}}$ is the sub-matrix of K^{-1} whose rows/columns are indexed by black/white vertices of \mathcal{E} .

- ▶ G infinite: Boltzmann measure \rightsquigarrow Gibbs measure
 - Periodic case [Cohn-Kenyon-Propp'01], [Ke.-Ok.-Sh.'06]
 - Non-periodic [dT'07], [Boutillier-dT'10], [B-dT-Raschel'19]

DIMER MODEL: PERIODIC CASE


- ▶ Assume G is bipartite, infinite, \mathbb{Z}^2 -periodic.



- ▶ Exhaustion of G by toroidal graphs: $(G_n) = (G/n\mathbb{Z}^2)$.

DIMER MODEL: PERIODIC CASE

- ▶ Fundamental domain: G_1




- ▶ Let K_1 be the Kasteleyn matrix of fundamental domain G_1 .
- ▶ Multiply edge-weights by $z, z^{-1}, w, w^{-1} \rightarrow K_1(z, w)$.
- ▶ The characteristic polynomial is:

$$P(z, w) = \det K_1(z, w).$$

Example: weight function $\nu \equiv 1$, $P(z, w) = 5 - z - \frac{1}{z} - w - \frac{1}{w}$.

DIMER MODEL: PERIODIC CASE

- Fundamental domain: G_1



- Let K_1 be the Kasteleyn matrix of **fundamental domain** G_1 .
- Multiply edge-weights by $z, z^{-1}, w, w^{-1} \rightarrow K_1(z, w)$.
- The **characteristic polynomial** is:

$$P(z, w) = \det K_1(z, w).$$


Example: weight function $\nu \equiv 1$, $P(z, w) = 5 - z - \frac{1}{z} - w - \frac{1}{w}$.

DIMER MODEL: SPECTRAL CURVE

- ▶ The spectral curve:

$$\mathcal{C} = \{(z, w) \in (\mathbb{C}^*)^2 : P(z, w) = 0\}.$$

- ▶ Amoeba: image of \mathcal{C} through the map $(z, w) \mapsto (\log |z|, \log |w|)$.



Amoeba of the square-octagon graph

DIMER MODEL AND HARNACK CURVES

THEOREMS

- ▶ Spectral curves of bipartite dimers

[Ke.-Ok.-Sh.'06] [Ke.-Ok.'06]
 \longleftrightarrow Harnack curves with points on ovals.

- ▶ Spectral curves of *isoradial*, bipartite dimer models with *critical weights* [Kenyon '02] [Kenyon-Okounkov'06]
 \longleftrightarrow Harnack curves of *genus 0*.

- ▶ Spectral curves of *minimal*, bipartite dimers [Goncharov-Kenyon '13]
 \longleftrightarrow Harnack curves with points on ovals.

Explicit (\longrightarrow) map

- ▶ [Fock'15] *Explicit* (\longleftarrow) map for all algebraic curves.
(no check on positivity).

DIMER MODEL AND HARNACK CURVES OF GENUS 1

THEOREM ([BOUTILLIER-DT-CIMASONI'19+])


Spectral curves of minimal, bipartite dimer models with Fock's weights

\longleftrightarrow

Harnack curves of genus 1 with a point on the oval.


QUAD-GRAPH, TRAIN-TRACKS

- ▶ Infinite, planar, embedded graph G ; embedded dual graph G^* .
- ▶ Corresponding **quad-graph** G^\diamond , **train-tracks**.




QUAD-GRAPH, TRAIN-TRACKS

- ▶ Infinite, planar, embedded graph G ; embedded dual graph G^* .
- ▶ Corresponding **quad-graph** G^\diamond , **train-tracks**.




QUAD-GRAPH, TRAIN-TRACKS

- ▶ Infinite, planar, embedded graph G ; embedded dual graph G^* .
- ▶ Corresponding **quad-graph** G^\diamond , **train-tracks**.




QUAD-GRAPH, TRAIN-TRACKS

- ▶ Infinite, planar, embedded graph G ; embedded dual graph G^* .
- ▶ Corresponding **quad-graph** G^\diamond , **train-tracks**.




QUAD-GRAPH, TRAIN-TRACKS

- ▶ Infinite, planar, embedded graph G ; embedded dual graph G^* .
- ▶ Corresponding **quad-graph** G^\diamond , **train-tracks**.



QUAD-GRAPH, TRAIN-TRACKS

- ▶ Infinite, planar, embedded graph G ; embedded dual graph G^* .
- ▶ Corresponding **quad-graph** G^\diamond , **train-tracks**.




ISORADIAL GRAPHS


- ▶ An **isoradial embedding** of an infinite, planar graph G is an embedding such that all of its faces are inscribed in a circle of radius 1, and such that the center of the circles are in the interior of the faces [Duffin] [Mercat] [Kenyon].
- ▶ Equivalent to: the quad-graph G^\diamond is embedded so that all its faces are rhombi.

THEOREM (KENYON-SCHLENCKER'04)


An infinite planar graph G has an isoradial embedding iff




ISORADIAL EMBEDDINGS



ISORADIAL EMBEDDINGS




ISORADIAL EMBEDDINGS




MINIMAL GRAPHS

- ▶ If the graph G is bipartite, train-tracks are naturally **oriented** (white vertex of the left, black on the right).



MINIMAL GRAPHS


- ▶ If the graph G is bipartite, train-tracks are naturally **oriented** (white vertex of the left, black on the right).
- ▶ A bipartite, planar graph G is **minimal** if



[Thurston'04] [Gulotta'08] [Ishii-Ueda'11] [Goncharov-Kenyon'13]

IMMERSIONS OF MINIMAL GRAPHS

- ▶ A **minimal isoradial immersion** of an infinite planar graph G is an immersion of the quadgraph G^\diamond such that:
 - all of the faces are rhombi (flat or reversed)



- the immersion is **flat**: the sum of the rhombus angles around every vertex and every face is equal to 2π .

PROPOSITION (BOUTILLIER-DT-CIMASONI'19+)

The flatness condition is equivalent to :

- *around every vertex there is at most one reversed rhombus*
- *around every face, the cyclic order of the vertices differ by at most disjoint transpositions in the embedding and in the immersion.*

THEOREM (BOUTILLIER-DT-CIMASONI'19+)

An infinite, planar, bipartite graph G has a minimal isoradial immersion iff it is minimal.

DIMER VERSION OF FOCK'S WEIGHTS

► Tool 1. Jacobi's (first) theta function.

- Parameter $q = e^{i\pi\tau}$, $\Im(\tau) > 0$, $\Lambda(q) = \pi\mathbb{Z} + \pi\tau\mathbb{Z}$, $\mathbb{T}(q) = \mathbb{C}/\Lambda$.

$$\theta(z) = 2q^{\frac{1}{4}} \sum_{n=0}^{\infty} (-1)^n q^{n(n+1)} \sin(2n+1)z.$$


- Allows to represent $\Lambda(q)$ -periodic meromorphic functions.
- $\theta(z) \sim 2q^{\frac{1}{4}} \sin(z)$ as $q \rightarrow 0$.

► Tool 2. Isoradially immersed, bipartite, minimal graph G .

- each train-track T is assigned **direction** $e^{i2\alpha_T}$.
- each edge $e = wb$ is assigned train-track directions $e^{2i\alpha}$, $e^{2i\beta}$, and a **half-angle** $\beta - \alpha \in [0, \pi)$.

DIMER VERSION OF FOCK'S ADJACENCY MATRIX

- **Tool 3. Discrete Abel map** [Fock] $D \in (\mathbb{R}/\pi\mathbb{Z})^{V(G^\circ)}$
 - Fix face f_0 and set $D(f_0) = 0$,
 - \circ : degree -1, \bullet : degree 1, faces: degree 0,
 - when crossing T : increase/decrease D by α_T accordingly.



- Point $t \in \frac{\pi}{2}\tau + \mathbb{R}$.
- **Fock's adjacency matrix**

$$K_{w,b}^{(t)} = \begin{cases} \frac{\theta(\beta - \alpha)}{\theta(t + D(b) - \beta)\theta(t + D(w) - \alpha)} & \text{if } w \sim b \\ 0 & \text{otherwise.} \end{cases}$$

DIMER VERSION OF FOCK'S ADJACENCY MATRIX

LEMMA ([BOUTILLIER-DT-CIMASONI'19+])

*Under the above assumptions, the matrix $K^{(t)}$ is a **Kasteleyn matrix** for a dimer model (positive weights) on G .*


FUNCTIONS IN THE KERNEL OF $K^{(t)}$

► Define $g^{(t)} : V^\diamond \times V^\diamond \times \mathbb{C} \rightarrow \mathbb{C}$

- $g_{x,x}^{(t)}(u) = 0,$
- If $f \sim w$, $g_{f,w}^{(t)}(u) = g_{w,f}^{(t)}(u)^{-1} = \frac{\theta(u + t + D(w))}{\theta(u - \alpha)},$
- If $f \sim b$, $g_{b,f}^{(t)}(u) = g_{f,b}^{(t)}(u)^{-1} = \frac{\theta(u - t - D(b))}{\theta(u - \alpha)},$

where $e^{2i\alpha}$ is the direction of the tt crossing the edge.

- If distance ≥ 2 , take product along path in G^\diamond .



PROPERTY OF THE FUNCTION $g^{(t)}$

LEMMA ([FOCK'15] [BOUTILLIER-DT-CIMASONI'19+])

- The function $g^{(t)}$ is well defined.
- The function $g^{(t)}$ is in the kernel of $K^{(t)}$:

$$\forall w \in W, x \in V^\diamond, \sum_{b:b \sim w} K_{w,b}^{(t)} g_{b,x}^{(t)}(u) = 0.$$

PROOF.

Weierstrass identity: $s, t \in \mathbb{T}(q)$, $a, b, c \in \mathbb{C}$,

$$\begin{aligned} & \frac{\theta(b-a)}{\theta(s-a)\theta(s-b)} \frac{\theta(u+s-a-b)}{\theta(u-a)\theta(u-b)} + \frac{\theta(c-b)}{\theta(s-b)\theta(s-c)} \frac{\theta(u+s-b-c)}{\theta(u-b)\theta(u-c)} + \\ & + \frac{\theta(a-c)}{\theta(s-c)\theta(s-a)} \frac{\theta(u+s-c-a)}{\theta(u-c)\theta(u-a)} = 0. \end{aligned}$$

□


EXPLICIT PARAMETERIZATION OF THE SPECTRAL CURVE

- ▶ Assume G is \mathbb{Z}^2 -periodic. Define the map ψ ,

$$\psi : \mathbb{T}(q) \rightarrow \mathbb{C}^2$$

$$u \mapsto \psi(u) = (z(u), w(u))$$

where $z(u) = g_{b_0, b_0+(1,0)}^{(t)}(u)$, $w(u) = g_{b_0, b_0+(0,1)}^{(t)}(u)$.




EXPLICIT PARAMETERIZATION OF THE SPECTRAL CURVE

PROPOSITION ([B-dT-C'19+])

The map ψ is an explicit birational parameterization of the spectral curve \mathcal{C} of the dimer model with Kasteleyn matrix $K^{(t)}$.

The real locus of \mathcal{C} is the image under ψ of the set $\mathbb{R}/\pi\mathbb{Z} \times \{0, \frac{\pi}{2}\tau\}$, where the connected component with ordinate $\frac{\pi}{2}\tau$ is bounded and the other is not.



(The spectral curve is independent of t).



GIBBS MEASURES FOR BIPARTITE DIMER MODELS

THEOREMS (KENYON-OKOUNKOV-SHEFFIELD'06)

- The dimer model on a \mathbb{Z}^2 -periodic, bipartite graph has a *two-parameter family* of ergodic Gibbs measures indexed by the *slope* (h, v) , i.e., by the average horizontal/vertical height change.
- The latter are obtained as weak limits of Boltzmann measures with *magnetic field coordinates* (B_x, B_y) .
- The *phase diagram* is given by the *amoeba* of the spectral curve \mathcal{C} .



LOCAL EXPRESSION FOR GIBBS MEASURES, GENUS 1

Suppose t fixed. Omit it from the notation.

THEOREM (BOUTILLIER-DT-CIMASONI'19+)

The 2-parameter set of EGM of the dimer model with Kasteleyn matrix K is $(\mathbb{P}^{u_0})_{u_0 \in D}$, where \forall subset of edges $\mathcal{E} = \{e_1 = w_1 b_1, \dots, e_n = w_n b_n\}$,

$$\mathbb{P}^{u_0}(e_1, \dots, e_n) = \left(\prod_{j=1}^n K_{w_j, b_j} \right) \det(A^{u_0})_{\mathcal{E}},$$


where $\forall b \in B, w \in W, A_{b,w}^{u_0} = \frac{i\theta'(0)}{2\pi} \int_{C_{b,w}^{u_0}} g_{b,w}(u) du$.

Moreover, when u_0


- is the unique point corresponding to the top boundary of D , the dimer model is *gaseous*,
- is in the interior of D , the dimer model is *liquid*,
- is a point corresponding to a cc of the lower boundary, the model is *solid*.

LOCAL EXPRESSIONS FOR ERGODIC GIBBS MEASURES, GENUS 1

► Domain D .



► Contours of integration.



COROLLARY

The slope of the Gibbs measure \mathbb{P}^{u_0} is:

$$h^{u_0} = \frac{1}{2\pi i} \int_{C^{u_0}} \frac{d}{du} (\log w(u)) du, \quad v^{u_0} = \frac{1}{2\pi i} \int_{C^{u_0}} \frac{d}{du} (\log z(u)) du.$$

IDEA OF THE PROOF

- **Proof 1.** Using [C-K-P], [K-O-S] the Gibbs measure \mathbb{P}^B with magnetic field coordinates $B = (B_x, B_y)$ has the following expression on cylinder sets:

$$\mathbb{P}^{(B_x, B_y)}(e_1, \dots, e_k) = \left(\prod_{j=1}^k K_{w_j, b_j} \right) \det(A^B)_\varepsilon,$$

where

$$A_{b+(m,n),w}^B = \int_{\mathbb{T}_B} \frac{Q(z, w)_{b,w}}{P(z, w)} z^{-m} w^{-n} \frac{dw}{2i\pi w} \frac{dz}{2i\pi z},$$

- Perform one integral by residues.
- Do the change of variable $u \mapsto \psi(u) = (z(u), w(u))$.
- Non-trivial cancellation !

IDEA OF THE PROOF

- ▶ **Proof 2.** Show that for every u_0 , A^{u_0} is an inverse of K .
 - Use Weierstrass identity.
 - Show that the contours of integration are such that one has 1 on the diagonal.

Use uniqueness statements of inverse operators.

CONSEQUENCES

- ▶ Suitable for asymptotics.
- ▶ Explicit local expressions for edge probabilities.

CONNECTION TO PREVIOUS WORK

- ▶ Genus 0. [Kenyon'02].
- ▶ Genus 1. Two specific cases were handled before:
 - the bipartite graph arising from the **Ising model** [Boutillier-dT-Raschel'19]
 - the $Z^{(t)}$ -**Dirac operator** [dT'18] \rightsquigarrow massive discrete holomorphic functions.

PERSPECTIVES

- ▶ 2-parameter family of Gibbs measures for **non-periodic graphs**.
Missing: every finite, simply connected subgraph of an isoradial immersion can be embedded in a **bipartite**, \mathbb{Z}^2 -periodic isoradial immersion.
- ▶ Extension to genus $g > 1$.
 - [Fock] gives a candidate for the dimer model.
 - Weierstrass identity \rightsquigarrow **Fay's trisecant identity**.