



Primordial Decoherence & Reliable Late-time Predictions



V. Vennin



R. Holman

2211.11046
2306.xxxxx



J. Martin



G. Kaplanek

Overview

Many QG puzzles are to do with horizons:
Black hole information loss; eternal inflation;...

Overview

Many QG puzzles are to do with horizons:
Black hole information loss; eternal inflation;...

Natural description using Open Quantum Systems
These can interestingly differ from Wilsonian description

Overview

Many QG puzzles are to do with late times:
Black hole information loss; eternal inflation;...

Overview

Many QG puzzles are to do with late times:
Black hole information loss; eternal inflation;...

Perturbative methods usually fail at late times
Inferences based on nearly free toy systems can be suspect

Overview

Many QG puzzles are to do with late times:
Black hole information loss; eternal inflation;...

Perturbative methods usually fail at late times
Inferences based on nearly free toy systems can be suspect

Open EFT tools can also allow predictions at late times
Decoherence of primordial fluctuations as an example...

Overview

In expanding universe Hubble length sets natural upper scale on correlations


$$-\Delta \phi = \ddot{\phi} + 3H\dot{\phi} + \frac{k^2}{a^2}\phi = 0$$

$$\text{with } H = \frac{\dot{a}}{a}$$

Modes are overdamped and freeze when $k/a \ll H$

Overview


Looking back into the past reveals ever more distant & disconnected Hubble patches



Overview


Primordial correlations require explanation

Leading explanations require quantum/gravity interplay




Overview

Correlations are a problem because relevant scales in past were super- Hubble



Overview

Solutions (eg inflation or bounces) change the naïve extrapolation




Guth 81, Linde 82, Albrecht & Steinhardt 82

If H is constant then $a(t) = e^{Ht}$

Overview

eg: inflation flattens out classical perturbations but constantly generates new quantum ones




Mukhanov & Chibisov 81, Guth & Pi 82, Starobinsky 82, Hawking 82

Resulting spectrum of fluctuations describes well the observed structure at later times

Overview

eg: inflation flattens out classical perturbations but constantly generates new quantum ones



Why do these initially quantum fluctuations look classical when seen at late times?

Decoherence

Why does this: $\langle \varphi_1 | \rho | \varphi_2 \rangle = \Psi[\varphi_1] \Psi^*[\varphi_2]$

Turn into this: $P[\varphi] = \langle \varphi | \rho | \varphi \rangle = |\Psi[\varphi]|^2$

Decoherence

Why does this: $\langle \varphi_1 | \rho | \varphi_2 \rangle = \Psi[\varphi_1] \Psi^*[\varphi_2]$

Turn into this: $P[\varphi] = \langle \varphi | \rho | \varphi \rangle = |\Psi[\varphi]|^2$

The subject of late-time (stochastic) cosmology

Starobinsky 86; Salopek & Bond 91; Starobinsky & Yokoyama 94;
CPB, Holman & Tasinato 16; Baumgart & Sundrum 19;
Gorbenko & Senatore 19; Mirbabayi 20, Cohen & Green 20

$$\partial_t P = \frac{H^3}{8\pi^2} \left(\frac{\partial^2 P}{\partial \varphi^2} \right) + \frac{\partial}{\partial \varphi} \left(\frac{\partial V}{\partial \varphi} P \right)$$

Decoherence

Why does this: $\langle \varphi_1 | \rho | \varphi_2 \rangle = \Psi[\varphi_1] \Psi^*[\varphi_2]$

Turn into this: $P[\varphi] = \langle \varphi | \rho | \varphi \rangle = |\Psi[\varphi]|^2$

*Does the density matrix become diagonal?
If so, why is the field basis the 'pointer' basis?*

$$\langle \varphi_1 | \rho | \varphi_2 \rangle \rightarrow |\Psi[\varphi_1]|^2 \delta[\varphi_1 - \varphi_2]$$

Decoherence

Why does this: $\langle \varphi_1 | \rho | \varphi_2 \rangle = \Psi[\varphi_1] \Psi^*[\varphi_2]$

Turn into this: $P[\varphi] = \langle \varphi | \rho | \varphi \rangle = |\Psi[\varphi]|^2$

*Does this question really need answering?
(Decoherence w/o decoherence)*

$$\langle F[\phi] \rangle = \text{Tr} (F \rho) = \int \mathcal{D}\varphi P[\varphi] F[\varphi]$$

Decoherence

Does this question really need answering (part 2)?

State squeezing makes super-Hubble modes 'WKB-classical'

Albrecht, et al 93, Starobinsky & Polarski 95

$$\hat{p} e^{\lambda S[\varphi]} = -i\lambda S'[\varphi] e^{\lambda S[\varphi]}$$

so $[\hat{p}, \hat{\phi}]$ is subdominant in $1/\lambda$ [or $k/(aH)$]

Decoherence

Even if not required, evolution of density matrix can be computed

Decoherence

Even if not required, evolution of density matrix can be computed

If decoherence occurs before horizon re-entry then quantum effects are unlikely to survive to be measured in the visible sky

Decoherence

Even if not required, evolution of density matrix can be computed

If decoherence occurs before horizon re-entry then quantum effects are unlikely to survive to be measured in the visible sky

Many have sought different sources for decoherence (other fields, thermal effects, stochastic evolution, etc)

Sakagami 88; Grischuk & Sidorov 89;

Brandenberger et al 90

Calzetta & Hu 95; Lesgourges et al 97; Kiefer et al 98;

...

Lombardo & Nacir 05; CPB Holman & Hoover 06;

Sharman & Moore 07; Kiefer et al 07; Kocksma &


Prokopec 11; CPB, Holman, Tasinato & Williams 14;

Nelson 16

...

Overview

Here argue that gravitational interactions amongst metric modes during inflation suffice to rapidly diagonalize super-Hubble density matrix in field basis



$$\delta_k(t) \sim \frac{\epsilon H^2}{M_p^2} \left(\frac{H a}{k} \right)^3 \propto e^{3Ht}$$

Outline

Open EFTs

Late-time control

Outline

Open EFTs

Late-time control

Cosmic decoherence

Metric fluctuations

& gravitational self-interactions




Open EFTs

Late-time control

Late-time perturbative breakdown

Perturbative methods generically break down at late times
(except possibly for scattering problems)

$$\exp\left[-i(H_0 + H_{\text{int}})t\right] \quad \text{vs} \quad e^{-iH_0 t} \left(1 - iH_{\text{int}}t + \dots\right)$$



Late-time perturbative breakdown

Perturbative methods generically break down at late times
(also happens in cosmology)

$$\mathcal{L} = -\sqrt{-g} \left[(\partial\phi)^2 + \frac{\lambda}{4!} \phi^4 \right]$$

$$\langle \phi^{2n} \rangle = (2n - 1)!! \left(\frac{H^2}{4\pi^2} \ln a \right)^n \left[1 - \frac{n(n+1)}{2} \left(\frac{\lambda}{36\pi^2} \right) \ln^2 a + \dots \right]$$

Late-time perturbative breakdown

Perturbative methods generically break down at late times
(also happens in cosmology)

$$\mathcal{L} = -\sqrt{-g} \left[(\partial\phi)^2 + \frac{\lambda}{4!} \phi^4 \right]$$

$$\langle \phi^{2n} \rangle = (2n - 1)!! \left(\frac{H^2}{4\pi^2} \ln a \right)^n \left[1 - \frac{n(n+1)}{2} \left(\frac{\lambda}{36\pi^2} \ln^2 a \right) + \dots \right]$$

$$a/a_0 = e^{H(t-t_0)}$$

Tsamis & Woodard 05

Late-time perturbative breakdown

Perturbative methods generically break down at late times
(also happens in cosmology)

$$\langle \phi^{2n} \rangle = (2n - 1)!! \left(\frac{H^2}{4\pi^2} \ln a \right)^n \left[1 - \frac{n(n+1)}{2} \left(\frac{\lambda}{36\pi^2} \right) \ln^2 a + \dots \right]$$

Tsamis & Woodard 05

$$\langle F[\phi] \rangle = \int \mathcal{D}\varphi P[\varphi] F[\varphi]$$

$$\partial_t P = \frac{H^3}{8\pi^2} \left(\frac{\partial^2 P}{\partial \varphi^2} \right) + \frac{\partial}{\partial \varphi} \left(\frac{\lambda \varphi^3}{3!} P \right)$$

Starobinsky 86

Starobinsky & Yokoyama 94

Stochastic inflation captures time evolution of IR sensitive correlations

Late-time perturbative breakdown

Perturbative methods generically break down at late times
(also happens in cosmology)

$$\langle \phi^{2n} \rangle = (2n - 1)!! \left(\frac{H^2}{4\pi^2} \ln a \right)^n \left[1 - \frac{n(n+1)}{2} \left(\frac{\lambda}{36\pi^2} \right) \ln^2 a + \dots \right]$$

Tsamis & Woodard 05

$$\partial_t P = \frac{H^3}{8\pi^2} \left(\frac{\partial^2 P}{\partial \varphi^2} \right) + \frac{\partial}{\partial \varphi} \left(\frac{\lambda \varphi^3}{3!} P \right)$$

$$P_\infty(\varphi) = P_0 \exp \left(-\frac{8\pi^2 V(\varphi)}{3} \right)$$

In this case secular growth signals non-Gaussian late-time distributions.

Late-time perturbative breakdown

Pedantic version of resummation argument


$$n(t) = n_0 e^{-\Gamma t} \quad \text{vs} \quad n(t) = n_0 - \Gamma n_0 t + \dots$$

$$\text{with } \Gamma \simeq \mathcal{O}(g^2)$$

Late-time perturbative breakdown

Pedantic version of resummation argument


$$n(t) \simeq n(t_0) \left[1 - \Gamma (t - t_0) + \dots \right]$$



Late-time perturbative breakdown

Pedantic version of resummation argument


$$n(t) \simeq n(t_1) \left[1 - \Gamma (t - t_1) + \dots \right]$$



Late-time perturbative breakdown

Pedantic version of resummation argument

$$\frac{\partial n}{\partial t} = -\Gamma n$$




Late-time perturbative breakdown

Integrating differential evolution resums all orders in $g^2 t$

$$n(t) = n_0 e^{-\Gamma t} \left[1 + \mathcal{O}(g^4 t) \right]$$


$$\text{with } \Gamma \simeq \mathcal{O}(g^2)$$



Late-time perturbative breakdown

Would NOT have worked if e.g. the differential eq depends on t_0

$$\frac{\partial n}{\partial t} = \int_{t_0}^t ds \Gamma(s) n(t - s)$$




Open systems

Wish to compute evolution when observations are restricted to a subsystem

$$H = H_{\text{sys}} + H_{\text{env}} + H_{\text{int}}$$

$$\rho_{\text{sys}}(t) = \text{Tr}_{\text{env}} \left[\rho_{\text{tot}}(t) \right]$$

$$\frac{\partial \rho_{\text{tot}}}{\partial t} = -i \left[H, \rho_{\text{tot}}(t) \right]$$




Open systems

Wish to compute evolution when observations are restricted to a subsystem

$$H = H_{\text{sys}} + H_{\text{env}} + H_{\text{int}}$$

$$\rho_{\text{sys}}(t) = \text{Tr}_{\text{env}} \left[\rho_{\text{tot}}(t) \right]$$

$$\frac{\partial \rho_{\text{sys}}}{\partial t} = -i \text{Tr}_{\text{env}} \left[H, \rho_{\text{tot}}(t) \right]$$



gfycat.com

Problem: RHS of evolution equation not expressed in terms of ρ_{sys} only

Open systems

Can very generally eliminate unobserved sector to obtain evolution for the system state in terms of environmental correlations

(Nakajima – Zwanzig equation)

Nakajima 58, Zwanzig 60

e.g. if $H_{\text{int}}(t) = A(t) \otimes B(t)$

$$\begin{aligned} \partial_t \rho_{\text{sys}} = & -i \left[A, \rho_{\text{sys}}(t) \right] \langle B \rangle_{\text{env}} \\ & + \int_{t_0}^t ds \left\{ C(s, t) \left[A(t), \rho_{\text{sys}}(s) A(s) \right] - C(t, s) \left[A(t), A(s) \rho_{\text{sys}}(s) \right] \right\} \\ & + \dots \end{aligned}$$

where $C(s, t) := \langle \delta B(s) \delta B(t) \rangle_{\text{env}}$

Open systems

Starting at second order the convolution over time makes NZ equation not generically useful for late-time resummation

$$\begin{aligned} \partial_t \rho_{\text{sys}}(t) = & -i [A, \rho_{\text{sys}}(t)] \langle B \rangle_{\text{env}} \\ & + \int_{t_0}^t ds \left\{ C(s, t) [A(t), \rho_{\text{sys}}(s) A(s)] - C(t, s) [A(t), A(s) \rho_{\text{sys}}(s)] \right\} \\ & + \dots \end{aligned}$$

where $C(s, t) := \langle \delta B(s) \delta B(t) \rangle_{\text{env}}$

Open systems

Starting at second order the convolution over time makes NZ equation not generically useful for late-time resummation

$$\begin{aligned} \partial_t \rho_{\text{sys}}(t) = & -i \left[A, \rho_{\text{sys}}(t) \right] \langle B \rangle_{\text{env}} \\ & + \int_{t_0}^t ds \left\{ C(s, t) \left[A(t), \rho_{\text{sys}}(s) A(s) \right] - C(t, s) \left[A(t), A(s) \rho_{\text{sys}}(s) \right] \right\} \\ & + \dots \end{aligned}$$

where $C(s, t) := \langle \delta B(s) \delta B(t) \rangle_{\text{env}}$

EFT PART: If C is sharply peaked at $s = t$ with width τ and if rest of integrand varies slowly compared to τ then can Taylor expand around $s = t$

Open systems

When such hierarchies exist then evolution becomes Markovian
(*Lindblad equation*)

Lindblad 76, Gorini et al 78

$$\begin{aligned} \partial_t \rho_{\text{sys}}(t) \simeq & -i \left[A, \rho_{\text{sys}}(t) \right] \langle B \rangle_{\text{env}} \\ & + \left\{ F(t_0, t) \left[A(t), \rho_{\text{sys}}(t) A(t) \right] - F(t, t_0) \left[A(t), A(t) \rho_{\text{sys}}(t) \right] \right\} \\ & + \dots \end{aligned}$$

where $F(t, t_0) := \int_{t_0}^t ds C(s, t)$

Open systems

When such hierarchies exist then evolution becomes Markovian

$$\begin{aligned} \partial_t \rho_{\text{sys}}(t) \simeq & -i [A, \rho_{\text{sys}}(t)] \langle B \rangle_{\text{env}} \\ & + \left\{ F(t_0, t) [A(t), \rho_{\text{sys}}(t) A(t)] - F(t, t_0) [A(t), A(t) \rho_{\text{sys}}(t)] \right\} \\ & + \dots \end{aligned}$$

$$\text{where } F(t, t_0) := \int_{t_0}^t ds C(s, t)$$

This can be useful for resummation if F is independent of t_0

Relevance for decoherence

First order term predicts Liouville evolution and so never contributes to decoherence

$$\partial_t \rho_{\text{sys}}(t) \simeq -i [A, \rho_{\text{sys}}(t)] \langle B \rangle_{\text{env}}$$

Relevance for decoherence

Second order term inequivalent to Hamiltonian evolution

$$+ \left\{ F(t_0, t) \left[A(t), \rho_{\text{sys}}(t) A(t) \right] - F(t, t_0) \left[A(t), A(t) \rho_{\text{sys}}(t) \right] \right\}$$

Relevance for decoherence

Second order term inequivalent to Hamiltonian evolution

$$+ \left\{ F(t_0, t) \left[A(t), \rho_{\text{sys}}(t) A(t) \right] - F(t, t_0) \left[A(t), A(t) \rho_{\text{sys}}(t) \right] \right\}$$

Second order term is simplest in basis for which A is diagonal and tends to drive off-diagonal components of ρ to zero in this basis

$$\text{e.g. if } A|\alpha\rangle = \alpha|\alpha\rangle$$

$$\partial_t \langle \alpha_1 | \rho_{\text{sys}} | \alpha_2 \rangle \ni -F (\alpha_1 - \alpha_2)^2 \langle \alpha_1 | \rho_{\text{sys}} | \alpha_2 \rangle$$

Relevance for decoherence

Second order term inequivalent to Hamiltonian evolution

$$+ \left\{ F(t_0, t) \left[A(t), \rho_{\text{sys}}(t) A(t) \right] - F(t, t_0) \left[A(t), A(t) \rho_{\text{sys}}(t) \right] \right\}$$

Second order term is simplest in basis for which A is diagonal and tends to drive off-diagonal components of ρ to zero in this basis

$$\text{e.g. if } A|\alpha\rangle = \alpha|\alpha\rangle$$

$$\partial_t \langle \alpha_1 | \rho_{\text{sys}} | \alpha_2 \rangle \ni -F (\alpha_1 - \alpha_2)^2 \langle \alpha_1 | \rho_{\text{sys}} | \alpha_2 \rangle$$

During inflation squeezing of the state ensures it is the field basis that generically diagonalizes A

Relevance for decoherence

Dependence of decoherence on scale factor is determined if correlations are localized in space

e.g. if
$$H_{\text{int}}(t) = \int d^3x a^3 \mathcal{A}(x, t) \otimes \mathcal{B}(x, t)$$

and
$$\langle \delta \mathcal{B}(x, t) \delta \mathcal{B}(x', t') \rangle = \frac{U(x, t) \delta^3(x - x') \delta(t - t')}{a^3}$$

Relevance for decoherence

Dependence of decoherence on scale factor is determined if correlations are localized in space

e.g. if
$$H_{\text{int}}(t) = \int d^3x a^3 \mathcal{A}(x, t) \otimes \mathcal{B}(x, t)$$

and
$$\langle \delta \mathcal{B}(x, t) \delta \mathcal{B}(x', t') \rangle = \frac{U(x, t) \delta^3(x - x') \delta(t - t')}{a^3}$$

then
$$\langle \alpha_1 | \rho_{\text{sys}}(t) | \alpha_2 \rangle = \langle \alpha_1 | \rho_{\text{sys}}(t_0) | \alpha_2 \rangle e^{-\Gamma(t)}$$

where
$$\Gamma(t) = \int d^3x dt \left[\alpha_1(x) - \alpha_2(x) \right]^2 a^3 U(x, t)$$

has width σ with $\sigma^{-1} \propto a^3 U(x, t)$

Relevance for decoherence

e.g. if $\mathcal{A}(x) \propto n(x)$

then for thermal fluctuations

$$U = n^2 \kappa_T T$$

so width is inversely proportional to temperature

scale factor is
normalized in space

(x, t)

$(t - t')$

then $\langle \alpha_1 | \rho_{\text{sys}}(t) | \alpha_2 \rangle = \langle \alpha_1 | \rho_{\text{sys}}(t_0) | \alpha_2 \rangle e^{-\Gamma(t)}$

where $\Gamma(t) = \int d^3x dt \left[\alpha_1(x) - \alpha_2(x) \right]^2 a^3 U(x, t)$

has width σ with $\sigma^{-1} \propto a^3 U(x, t)$

A scenic view of a mountain range with a dense forest in the foreground and a clear blue sky. The mountains are rugged and rocky, with some snow patches. The forest is a mix of evergreen and deciduous trees. The sky is a clear, light blue with a few wispy clouds.

Cosmic Decoherence

*Lindblad for metric
fluctuations*

Single clock metric fluctuations

For simplest models a single field controls breaking of de Sitter symmetries near horizon exit

$$\mathcal{L} = -\sqrt{-g} \left[M_p^2 \mathcal{R} + (\partial\phi)^2 + V(\phi) \right]$$

$$\phi = \varphi(t)$$

and $ds^2 = a^2 \left[-d\eta^2 + \delta_{ij} dx^i dx^j \right]$

with $3M_p^2 H^2 = \rho = \frac{1}{2} \dot{\varphi}^2 + V(\varphi)$

Single clock metric fluctuations

For simplest models a single field controls breaking of de Sitter symmetries near horizon exit

$$\mathcal{L} = -\sqrt{-g} \left[M_p^2 \mathcal{R} + (\partial\phi)^2 + V(\phi) \right]$$

$$\phi = \varphi(t)$$

$$\text{and } ds^2 = a^2 \left[-d\eta^2 + \delta_{ij} dx^i dx^j \right]$$

$$\text{with } 3M_p^2 H^2 = \rho = \frac{1}{2} \dot{\varphi}^2 + V(\varphi)$$

$$\text{slow-roll parameter } \epsilon_1 = -\frac{\dot{H}}{H^2} \simeq \frac{1}{2} (M_p V' / V)^2 \ll 1$$

Single clock metric fluctuations

Primordial fluctuations are described by deviations from homogeneous background

$$\mathcal{L} = -\sqrt{-g} \left[M_p^2 \mathcal{R} + (\partial\phi)^2 + V(\phi) \right]$$

$$\phi = \varphi(t) + \delta\phi(x, t)$$

and $ds^2 = a^2 \left[-(1 + 2\psi)d\eta^2 + e^{2\zeta}(\delta_{ij} + h_{ij})dx^i dx^j \right]$

Single clock metric fluctuations

Primordial fluctuations are described by deviations from homogeneous background

$$\mathcal{L} = -\sqrt{-g} \left[M_p^2 \mathcal{R} + (\partial\phi)^2 + V(\phi) \right]$$

$$\phi = \varphi(t) + \delta\phi(x, t)$$

and $ds^2 = a^2 \left[-(1 + 2\psi)d\eta^2 + e^{2\zeta} (\delta_{ij} + h_{ij}) dx^i dx^j \right]$

semiclassical control parameter $(H^2 / 4\pi M_p^2) \ll 1$

Single clock metric fluctuations

Primordial fluctuations are described by deviations from homogeneous background

$$\mathcal{L} = -\sqrt{-g} \left[M_p^2 \mathcal{R} + (\partial\phi)^2 + V(\phi) \right]$$

$$\phi = \varphi(t) + \delta\phi(x, t)$$

and $ds^2 = a^2 \left[-(1 + 2\psi)d\eta^2 + e^{2\zeta} (\delta_{ij} + h_{ij}) dx^i dx^j \right]$


other useful variables

$$v = a \left(\delta\phi + \varphi' \psi / H \right) \quad \text{spatially flat gauge}$$

$$\zeta = (\delta\phi / \sqrt{\epsilon} M_p) + \dots \quad \text{comoving gauge}$$


System vs environment

Only a specific range of fluctuation modes is visible to late-time cosmologists



System vs environment

Define environment to be the unobserved (in particular shorter wavelength) modes



$$v = v_{\text{sys}}(k < k_*) + v_{\text{env}}(k > k_*)$$

System vs environment

Nonlinearity of GR implies many self interactions amongst metric fluctuations:

$$\mathcal{L}_{\text{int}} = \frac{\partial^2 h^3}{M_p} + \frac{\partial^2 h^4}{M_p^2} + \frac{\partial^2 h^5}{M_p^3} + \dots$$

Dominant contribution arises at order M_p^{-2} and so is 2nd order in cubic interactions or first order in quartic interactions

BUT first order never contributes to decoherence: *leaves only cubic terms*

System vs environment

Complete basis of cubic interactions amongst metric fluctuations is known

Maldacena 04

$$\mathcal{L}_{sss} = \epsilon^2 M_p^2 a(\partial\zeta)^2\zeta + \dots$$

$$\mathcal{L}_{stt} = \epsilon M_p^2 a(\partial_k h_{ij})^2\zeta + \dots$$

$$\mathcal{L}_{sst} = \epsilon M_p^2 a(\partial_i\zeta\partial_j\zeta)h^{ij} + \dots$$

plus subdominant terms (higher order in slow roll; $d\zeta/dt$ terms; ...)

System vs environment

Must divide these up into system and environmental parts, for example for

$$\mathcal{L} = \epsilon^2 M_p^2 a(\partial\zeta)^2\zeta + \dots$$

Only linear and cubic terms possible for long-wavelength sector

$$\mathcal{L} \ni v_{\text{sys}}(\partial v_{\text{sys}})^2, v_{\text{sys}}(\partial v_{\text{env}})^2, v_{\text{env}}(\partial v_{\text{env}} \cdot \partial v_{\text{sys}}), \\ v_{\text{env}}(\partial v_{\text{sys}})^2, v_{\text{env}}(\partial v_{\text{env}})^2, \dots$$

System vs environment

Must divide these up into system and environmental parts, for example for

$$\mathcal{L} = \epsilon^2 M_p^2 a(\partial\zeta)^2\zeta + \dots$$

Only linear and cubic terms possible for long-wavelength sector

$$\mathcal{L} \ni v_{\text{sys}}(\partial v_{\text{sys}})^2, v_{\text{sys}}(\partial v_{\text{env}})^2, v_{\text{env}}(\partial v_{\text{env}} \cdot \partial v_{\text{sys}}), \\ v_{\text{env}}(\partial v_{\text{sys}})^2, v_{\text{env}}(\partial v_{\text{env}})^2, \dots$$

Doesn't couple system to environment

System vs environment

Must divide these up into system and environmental parts, for example for

$$\mathcal{L} = \epsilon^2 M_p^2 a(\partial\zeta)^2\zeta + \dots$$

Only linear and cubic terms possible for long-wavelength sector

$$\mathcal{L} \ni v_{\text{sys}}(\partial v_{\text{sys}})^2, v_{\text{sys}}(\partial v_{\text{env}})^2, v_{\text{env}}(\partial v_{\text{env}} \cdot \partial v_{\text{sys}}), \\ v_{\text{env}}(\partial v_{\text{sys}})^2, v_{\text{env}}(\partial v_{\text{env}})^2, \dots$$

Doesn't couple system to environment
Forbidden by momentum conservation

System vs environment

Must divide these up into system and environmental parts, for example for

$$\mathcal{L} = \epsilon^2 M_p^2 a(\partial\zeta)^2\zeta + \dots$$

Only linear and cubic terms possible for long-wavelength sector

$$\mathcal{L} \ni v_{\text{sys}}(\partial v_{\text{sys}})^2, v_{\text{sys}}(\partial v_{\text{env}})^2, v_{\text{env}}(\partial v_{\text{env}} \cdot \partial v_{\text{sys}}), \\ v_{\text{env}}(\partial v_{\text{sys}})^2, v_{\text{env}}(\partial v_{\text{env}})^2, \dots$$

Doesn't couple system to environment

Forbidden by momentum conservation

Subdominant because derivatives are small

Gravitational Lindblad equation

Following general steps leads to Markovian evolution
for super-Hubble modes

$$\mathcal{L} = \epsilon^2 M_p^2 a(\partial\zeta)^2\zeta + \dots$$

Environment and system coupled to one another through

$$H_{\text{int}} = \int d^3x \frac{\sqrt{\epsilon}}{a M_p} v_{\text{sys}} \otimes \mathcal{B}_{\text{env}}$$

and so to second order evolution ***does not change mode label k*** and involves environmental correlations of

$$\mathcal{B}_{\text{env}} \propto \delta^{ij} \partial_i v_{\text{env}} \partial_j v_{\text{env}}$$

Gravitational Lindblad equation

Evolution of each mode is Markovian at late times for super-Hubble modes, with Schrodinger-picture Lindblad equation:

$$\frac{\mathcal{V}}{(2\pi)^3} \frac{\partial \rho_k}{\partial \eta} \simeq - \text{Re} F_k(\eta, \eta_{\text{in}}) \left[v_k, [v_k, \rho_k(\eta)] \right] \\ - i \left[\mathcal{H}_k^0(\eta) - \text{Im} F_k(\eta, \eta_{\text{in}}) v_k^2, \rho_k(\eta) \right]$$

Gravitational Lindblad equation

Evolution of each mode is Markovian at late times for super-Hubble modes, with Schrodinger-picture Lindblad equation:

$$\frac{\mathcal{V}}{(2\pi)^3} \frac{\partial \rho_k}{\partial \eta} \simeq - \text{Re} F_k(\eta, \eta_{\text{in}}) \left[v_k, [v_k, \rho_k(\eta)] \right] \\ - i \left[\mathcal{H}_k^0(\eta) - \underline{\text{Im} F_k(\eta, \eta_{\text{in}}) v_k^2}, \rho_k(\eta) \right]$$

where $\text{Im} F_k$ contains a **UV divergence** with the right form to be absorbed into the standard Einstein-Hilbert and curvature-squared counterterms

$$\text{Im} F_k(\eta, \eta_{\text{in}}) \ni \frac{\epsilon H^2 k^2}{1024 \pi^2 M_p^2} \left(\frac{40}{(-k\eta)^2} - \frac{92}{3} + \frac{43}{15} (-k\eta)^2 \right) \\ \times \left[\frac{2}{n-4} + \log \left(\frac{2k_* + k}{\mu} \right) \right]$$

Gravitational Lindblad equation

Evolution of each mode is Markovian at late times for super-Hubble modes, with Schrodinger-picture Lindblad equation:

$$\frac{\mathcal{V}}{(2\pi)^3} \frac{\partial \rho_k}{\partial \eta} \simeq - \underline{\text{Re}F_k(\eta, \eta_{\text{in}})} \left[v_k, [v_k, \rho_k(\eta)] \right] - i \left[\mathcal{H}_k^0(\eta) - \text{Im}F_k(\eta, \eta_{\text{in}}) v_k^2, \rho_k(\eta) \right]$$

Decoherence comes purely from $\text{Re} F_k$ which is **UV finite** and has a **universal form** (independent of k_* and η_{in}) at late super-Hubble times

$$\text{Re}F_k(\eta, \eta_{\text{in}}) \simeq \frac{\epsilon H^2 k^2}{1024\pi^2 M_p^2} (1 + 2) \left[\frac{20\pi}{(-k\eta)^2} + \frac{g(k_*/k, -k\eta_{\text{in}})}{(-k\eta)} + \dots \right]$$

Gravitational Lindblad equation

Evolution of each mode is Markovian at late times for super-Hubble modes, with Schrodinger-picture Lindblad equation:

$$\frac{\mathcal{V}}{(2\pi)^3} \frac{\partial \rho_k}{\partial \eta} \simeq - \text{Re} F_k(\eta, \eta_{\text{in}}) \left[v_k, [v_k, \rho_k(\eta)] \right] - i \left[\mathcal{H}_k^0(\eta) - \text{Im} F_k(\eta, \eta_{\text{in}}) v_k^2, \rho_k(\eta) \right]$$

Decoherence comes purely from $\text{Re} F_k$ which is **UV finite** and has a **universal form** (independent of k_* and η_{in}) at late super-Hubble times

$$\text{Re} F_k(\eta, \eta_{\text{in}}) \simeq \frac{\epsilon H^2 k^2}{1024 \pi^2 M_p^2} (1 + 2) \left[\frac{20\pi}{(-k\eta)^2} - \frac{g(k_*/k, -k\eta_{\text{in}})}{(-k\eta)} + \dots \right]$$

Gravitational Lindblad equation

Evolution of each mode is Markovian at late times for super-Hubble modes, with Schrodinger-picture Lindblad equation:

$$\frac{\mathcal{V}}{(2\pi)^3} \frac{\partial \rho_k}{\partial \eta} \simeq - \text{Re} F_k(\eta, \eta_{\text{in}}) \left[v_k, [v_k, \rho_k(\eta)] \right] - i \left[\mathcal{H}_k^0(\eta) - \text{Im} F_k(\eta, \eta_{\text{in}}) v_k^2, \rho_k(\eta) \right]$$

Decoherence comes purely from $\text{Re} F_k$ which is **UV finite** and has a **universal form** (independent of k_* and η_{in}) at late super-Hubble times

$$\text{Re} F_k(\eta, \eta_{\text{in}}) \simeq \frac{\epsilon H^2 k^2}{1024 \pi^2 M_p^2} (1 + 2) \left[\frac{20\pi}{(-k\eta)^2} + \frac{g(k_*/k, -k\eta_{\text{in}})}{(-k\eta)} + \dots \right]$$

Scalar environment contributes 1 while tensor environment contributes 2

Cosmic decoherence

Can quantify the decoherence for each mode using

$$\text{Tr} [\rho_k^2] = \frac{1}{\sqrt{1 + 4\delta_k(\eta)}}$$

Since the state is coherent if and only if $\delta_k = 0$

Cosmic decoherence

Can quantify the decoherence for each mode using

$$\text{Tr} [\rho_k^2] = \frac{1}{\sqrt{1 + 4\delta_k(\eta)}}$$

Since the state is coherent if and only if $\delta_k = 0$

Lindblad evolution makes the following late-time prediction

$$\delta_k(\eta) \simeq \frac{5(1 + 2)\epsilon H^2}{96\pi M_p^2 (-k\eta)^3} \simeq \frac{5\epsilon H^2}{32\pi M_p^2} \left(\frac{aH}{k} \right)^3$$

Cosmic decoherence

Can quantify the decoherence for each mode using

$$\text{Tr} [\rho_k^2] = \frac{1}{\sqrt{1 + 4\delta_k(\eta)}}$$

Since the state is coherent if and only if $\delta_k = 0$

Lindblad evolution makes the following late-time prediction

$$\delta_k(\eta) \simeq \frac{5(1 + 2)\epsilon H^2}{96\pi M_p^2 (-k\eta)^3} \simeq \frac{5\epsilon H^2}{32\pi M_p^2} \left(\frac{aH}{k}\right)^3$$

Starts small: $\epsilon^2 \left(\frac{H^2}{32\pi\epsilon M_p^2}\right) < 10^{-4} \times 10^{-10}$

Cosmic decoherence

Can quantify the decoherence for each mode using

$$\text{Tr} [\rho_k^2] = \frac{1}{\sqrt{1 + 4\delta_k(\eta)}}$$

Since the state is coherent if and only if $\delta_k = 0$

Lindblad evolution makes the following late-time prediction

$$\delta_k(\eta) \simeq \frac{5(1+2)\epsilon H^2}{96\pi M_p^2 (-k\eta)^3} \simeq \frac{5\epsilon H^2}{32\pi M_p^2} \left(\frac{aH}{k}\right)^3$$

Grows quickly: $\propto \exp(+3Ht)$

Cosmic decoherence

Can quantify the decoherence for each mode using

$$\text{Tr} [\rho_k^2] = \frac{1}{\sqrt{1 + 4\delta_k(\eta)}}$$

Since the state is coherent if and only if $\delta_k = 0$

Similar calculation gives effects of scalar environment on tensors

$$\delta_k(\eta) \simeq \frac{H^2}{72\pi M_p^2} \left(\frac{aH}{k} \right)^3$$

For tensor modes decoherence is not slow-roll suppressed



AUTRES
DIRECTIONS

TOUTES
DIRECTIONS

The way forward

Takeaway messages

Takeaway Messages

What decoheres primordial quantum fluctuations?

Do not yet know definitively, but Open EFT reasoning provides tools for deciding within any particular scenario

Takeaway Messages

What decoheres primordial quantum fluctuations?

Do not yet know definitively, but Open EFT reasoning provides tools for deciding within any particular scenario

Preliminary evidence is that unseen short wavelength modes have ample time to decohere observed fluctuations even if they only couple with gravitational strength as predicted by GR. Likely a floor for how quickly primordial fluctuations decohere.

Takeaway Messages

What decoheres primordial quantum fluctuations?

Do not yet know definitively, but Open EFT reasoning provides tools for deciding within any particular scenario

Preliminary evidence is that unseen short wavelength modes have ample time to decohere observed fluctuations even if they only couple with gravitational strength as predicted by GR. Likely a floor for how quickly primordial fluctuations decohere.

Are there observable consequences?

Possibly so, but quantum coherence effects only possible if observed fluctuations do not spend too much super-Hubble time during inflation



Takeaway Messages

Perturbative methods are usually unreliable at late times

It can be dangerous to extrapolate free-field behaviour arbitrarily far into the future since this is implicitly perturbative

Many tools exist elsewhere in physics to deal with this kind of situation and it behooves us to use them for gravitational settings too.

Takeaway Messages

Perturbative methods are usually unreliable at late times

It can be dangerous to extrapolate free-field behaviour arbitrarily far into the future since this is implicitly perturbative

Many tools exist elsewhere in physics to deal with this kind of situation and it behooves us to use them for gravitational settings too.

Some predictions can be controlled

Small things can accumulate to cause dramatic effects at very late times. These effects need not carry lots of energy or destabilize the geometry.





fin



fin