

# On a new proof of the Harris ergodic theorem and some applications

José A. Cañizo  
University of Granada

*Based on collaborations with **Chu Qi Cao** (Paris-Dauphine),  
**J. Evans** (U. Cambridge), **S. Mischler** (Paris-Dauphine),  
**H. Yoldaş** (BCAM & U. Granada)*

Banff, April 13 2018

# Convergence to equilibrium of Markov processes

Studying convergence to equilibrium of Markov processes is a central problem in many fields. Some examples we are interested in:

- Linear Boltzmann and related models
- PDEs in mathematical biology
- Integro-differential / Nonlocal PDE (fractional / nonlocal diffusion)
- Kinetic-type linear PDE, typically hypocoercive

We look at this problem from a PDE point of view and often we want to extend our arguments to:

- Nonlinear equations
- Non-conservative equations

Many ideas have been used to prove convergence to equilibrium of Markov processes:

**Entropy methods** Useful for several PDE in mathematical physics. *Hypocoercive* equations are an obstacle. *One of the main topics of this conference.*

**Operator splitting** Often gives a non-constructive spectral gap. “Weyl method”, classical in kinetic theory / operator theory. See also [**Gualdani, Mischler & Mouhot**].

**Doeblin / Harris / Meyn-Tweedie** The method we’re discussing today. Only applicable to linear problems.

- **[Harris 1956]**
- **[Meyn & Tweedie 1992, 1993]**
- **[Hairer & Mattingly 2011]** - Simplified proof using mass transport distances.
- **[Bakry, Cattiaux & Guillin 2008], [Douc, Fort & Guillin 2009]** - Subexponential version
- **[Gabriel 2017]** - Renewal equation
- **[Dumont & Gabriel 2017]** - Integrate-and-fire neuron model
- **[Eberle, Guillin & Zimmer 2016], [Hu & Wang 2017]** - Kinetic Fokker-Planck
- **[Bansaye, Cloez & Gabriel 2017]** - Doeblin theorem for non-conservative cases

# Simple statement: Doeblin's theorem

$(S_t)_{t \geq 0}$  Markov semigroup, where  $S_t$  represents the evolution of the law of the process (so  $S_t$  acts on measures). Assume:

## Doeblin's condition

There exists  $t_0 > 0$ ,  $0 < \alpha < 1$  and  $\nu \in \mathcal{P}$  with

$$S_{t_0}\mu \geq \alpha\nu \quad \text{for all } \mu \in \mathcal{P}. \quad (1)$$

Then there exists a unique equilibrium  $\mu_* \in \mathcal{P}$  and

$$\|S_t(\mu - \mu_*)\| \leq \frac{1}{1 - \alpha} e^{-\lambda t} \|n - n_*\|, \quad \text{for } t \geq 0 \quad (2)$$

for all  $\mu \in \mathcal{P}$ , where

$$\lambda := -\frac{\log(1 - \alpha)}{t_0} > 0.$$

# Doebli's theorem: proof

The result follows easily from this: first, **if**  $\mu_1, \mu_2 \in \mathcal{P}$  **have disjoint support**, by the triangle inequality we have

$$\|\mathcal{S}\mu_1 - \mathcal{S}\mu_2\|_{\text{TV}} \leq \|\mathcal{S}\mu_1 - \alpha\nu\|_{\text{TV}} + \|\mathcal{S}\mu_2 - \alpha\nu\|_{\text{TV}}.$$

Now, since  $\mathcal{S}\mu_i \geq \alpha\nu$ , we can write

$$\|\mathcal{S}\mu_i - \alpha\nu\|_{\text{TV}} = \int (\mathcal{S}\mu_i - \alpha\nu) = \int \mu_i - \alpha = 1 - \alpha,$$

which gives

$$\|\mathcal{S}\mu_1 - \mathcal{S}\mu_2\| \leq 2(1 - \alpha) = (1 - \alpha)\|\mu_1 - \mu_2\|.$$

To get this for any  $\mu_1, \mu_2 \in \mathcal{P}$ , write

$$\mu_1 - \mu_2 = (\mu_1 - \mu_2)_+ - (\mu_1 - \mu_2)_-$$

and apply previous case.

# Harris' theorem

## Lyapunov condition

There exists  $t_0 > 0$ ,  $0 < \gamma < 1$ ,  $K \geq 0$  and a measurable function  $V: \Omega \rightarrow [1, +\infty)$  such that

$$\int V S_{t_0} \mu \leq \gamma \int V \mu + K \int \mu \quad \text{for all nonnegative } \mu \in \mathcal{M}. \quad (3)$$

## Local Doeblin's condition

There exists  $0 < \alpha < 1$  and  $\nu \in \mathcal{P}$  with

$$S_{t_0} \mu \geq \alpha \nu \quad \text{for all } \mu \in \mathcal{P} \text{ supported on a "large" set } \mathcal{C} \quad (4)$$

Then there exists a unique equilibrium  $n_* \in \mathcal{P}$  and

$$\|S_t(\mu - \mu_*)\|_{\beta} \leq C e^{-\lambda t} \|\mu - \mu_*\|_{\beta}. \quad \text{for } t \geq 0 \quad (5)$$

for some  $\lambda > 0$ ,  $C \geq 1$ ,  $\beta > 0$  and all  $\mu \in \mathcal{P}$ .



# Example: kinetic linear BGK

We consider

$$\partial_t f + v \cdot \nabla_x f = Lf,$$

posed for  $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$ , where

$$Lf(x, v) := L^+ f(x, v) - f(x, v) := \left( \int_{\mathbb{R}^d} f(x, u) du \right) M(v) - f(x, v).$$

Call  $T_t \equiv$  transport semigroup associated to  $-v \nabla_x f$ . Then:

$$f_t \geq e^{-t} \int_0^t \int_0^s T_{t-s} L^+ T_{s-r} L^+ T_r f_0 dr ds.$$



# Two bounds

## Jumping to small velocities

There exist  $\alpha_L, \delta_L > 0$  such that for all  $x$ ,

$$L^+ \delta_{(x,v)} \geq \alpha_L \delta_x \mathbb{1}[|v| \leq \delta_L].$$

## Jumping to small positions

There exist  $\alpha_T, \delta_T, t_0, \epsilon$  such that for all  $x$ ,

$$\int_{\mathbb{R}^d} T_t \left( \delta_x \mathbb{1}[|v| \leq \delta_L] \right) dv \geq \alpha_T \mathbb{1}[|x| \leq \delta_T] \quad \forall t \in (t_0 - \epsilon, t_0 + \epsilon).$$

This allows us to use Doeblin's Theorem directly and obtain

$$\|f_t - M\|_1 \leq C e^{-\lambda t} \|f_0 - M\|_1.$$

# Example: kinetic linear BGK + potential

We consider

$$\partial_t f + v \cdot \nabla_x f - \nabla_x \Phi(x) \cdot \nabla_v f = Lf,$$

posed for  $(x, v) \in \mathbb{T}^d \times \mathbb{R}^d$ , where

$$Lf(x, v) := L^+ f(x, v) - f(x, v) := \left( \int_{\mathbb{R}^d} f(x, u) du \right) M(v) - f(x, v).$$

Call  $T_t \equiv$  semigroup associated to  $-v \nabla_x f + \nabla_x \Phi(x) \cdot \nabla_v f$ .

Then:

$$f_t \geq e^{-t} \int_0^t \int_0^s T_{t-s} L^+ T_{s-r} L^+ T_r f_0 dr ds.$$

# Two bounds (with potential $\Phi$ )

## Jumping to small velocities

There exist  $\alpha_L, \delta_L > 0$  such that

$$L^+ \delta_{(x,v)} \geq \alpha_L \delta_x \mathbb{1}[|v| \leq \delta_L].$$

## Jumping to small positions

Assume  $|\nabla\Phi(x)| \leq C|\Phi(x)|^\eta$  for some  $0 < \eta < 1$ . For any  $R_1$  there exists  $\delta_M, R_2, \epsilon, t_0$  such that

$$\int T_s(\delta_{x_0} \mathbb{1}[|v| \leq R_2]) dv \geq \mathbb{1}[|x| \leq \delta_M] \quad \text{for } t \in [t_0 - \epsilon, t_0 + \epsilon]$$

for any  $\Phi(x_0) \leq R_1$

We cannot obtain this for all  $x_0$ , so we need to use Harris instead of Doeblin.

The Foster-Lyapunov condition works with:

$$V(x, v) := 1 + \Phi(x) + \frac{1}{2}|v|^2 + \frac{1}{4}x \cdot v + \frac{1}{8}|x|^2$$

This allows us to use Harris' Theorem directly and obtain

$$\|f_t - M\|_* \leq Ce^{-\lambda t} \|f_0 - M\|_*$$

with

$$\|g\|_* := \int |g| + \int V|g|.$$

# Harris' theorem: proof

This proof is a simplification of that in **[Hairer & Mattingly 2011]**.

The norm  $\|\cdot\|_\beta$  is here

$$\|\mu\|_\beta := \int |\mu| + \beta \int |\mu| V$$

It is enough to show that

$$\|\mathcal{S}_{t_0}\mu\|_\beta \leq (1 - \delta)\|\mu\|_\beta$$

for all measures  $\mu$  with  $\int \mu = 0$ . (That is,  $\mathcal{S}_{t_0}$  is contractive in the  $\|\cdot\|_\beta$  norm). We extend our previous idea as follows:

# First case: Contractivity for large $V$ -moment.

Take  $\delta \in (0, 1 - \gamma)$ . If we have

$$\int V|\mu| > \frac{K}{1 - \gamma - \delta} \int |\mu| \quad (6)$$

then

$$\begin{aligned} \|S\mu\|_V &= \int V|S\mu| \leq \gamma \int V|\mu| + K \int |\mu| \\ &\leq (1 - \delta) \int V|\mu| = (1 - \delta) \|\mu\|_V. \end{aligned}$$

This easily implies the contractivity of the norm  $\|\cdot\|_\beta$  in this case.

## Second case: Contractivity for small $V$ -moment

Assume that

$$\int V|\mu| < \frac{K}{1-\gamma-\delta} \int |\mu|, \quad (7)$$

In this case a sizeable part of the mass of  $\mu_+$  and  $\mu_-$  is in  $\mathcal{C}$ :

$$\int_{\mathcal{C}} \mu_{\pm} \geq (1-\xi) \int \mu_{\pm}.$$

Using this we can carry out a similar argument as in Doeblin's Theorem: we have

$$\mathbf{S}\mu_{\pm} \geq \alpha\nu \int_{\mathcal{C}} \mu_{\pm} \geq \alpha(1-\xi)\nu \int \mu_{\pm} =: \eta$$

and then

$$\begin{aligned} \|\mathbf{S}\mu\| &\leq \|\mathbf{S}\mu_+ - \eta\| + \|\mathbf{S}\mu_- - \eta\| = \int \mathbf{S}\mu_+ + \int \mathbf{S}\mu_- - 2 \int \eta \\ &= \int \mu_+ + \int \mu_- - \alpha(1-\xi) \int |\mu| = (1 - \alpha(1-\xi)) \int |\mu|. \end{aligned}$$

It is then easy to complete the argument to obtain contractivity in the  $\|\cdot\|_\beta$  norm, for  $\beta$  small enough:

$$\begin{aligned}\|\mathbf{S}\mu\|_\beta &= \|\mathbf{S}\mu\| + \beta \int \mathbf{V}|\mathbf{S}\mu| \\ &\leq (1 - \alpha' + \beta K)\|\mu\| + \beta\gamma \int \mathbf{V}|\mu| \leq (1 - \delta_1)\|\mu\|_\beta,\end{aligned}$$

with

$$\delta_1 := \min\{\alpha' - \beta K, 1 - \gamma\}. \quad (8)$$

Taking then  $\beta K < \alpha'$  gives the contractivity in this case.



# Harris' theorem, subexponential version

$L \equiv$  generator of the semigroup.

## Lyapunov condition

There exists  $K \geq 0$  and a measurable function  $V: \Omega \rightarrow [1, +\infty)$  such that

$$\int VL\mu \leq - \int \varphi(V)\mu + K \int \mu \quad \text{for all } \mu \in \mathcal{P}.$$

## Local Doeblin's condition

There exists  $0 < \alpha < 1$  and  $\nu \in \mathcal{P}$  with

$$S_{t_0}\mu \geq \alpha\nu \quad \text{for all } \mu \in \mathcal{P} \text{ supported on a "large" set } \mathcal{C} \quad (9)$$

Then there exists a unique equilibrium  $n_* \in \mathcal{P}$  and

$$\|S_t(n - n_*)\| \leq r(t)\|n - n_*\|_V. \quad \text{for } t \geq 0 \quad (10)$$

for some  $r(t) \rightarrow 0$  as  $t \rightarrow +\infty$  and all  $n \in \mathcal{P}$ .



We can in fact take

$$r(t) := \frac{1}{\varphi(H^{-1}(t))}$$

where

$$H(v) := \int_1^v \frac{1}{\varphi(u)} du.$$

# Harris' theorem, subexponential version II

For the moment we can prove a weaker version with the same technique: we add the condition:

## Boundedness of moments

There exists  $V_2 = V_2(x)$  with  $V(x)/V_2(x) \rightarrow 0$  as  $x \rightarrow +\infty$  such that

$$\int |S_t \mu| V_2 \leq C \int |\mu| V_2 \quad \text{for all } t \geq 0, \mu \in \mathcal{P}.$$

Then we obtain the same conclusion with a different  $r(t)$ , which decays faster if  $V_2$  is larger.

# Subexponential version: proof

**First case.** *Estimate for large  $V$ -moment.* Assume that for  $t \in [t_n, t_n + T]$  we have

$$\int V_0 |\mu_t| \geq \frac{2K}{c_1} \int |\mu_t|.$$

Then

$$\frac{d}{dt} \int V_1 |\mu_t| \leq -K \int V_0 |\mu_t|.$$

Consequently, if we choose any  $\epsilon > 0$  we have

$$\begin{aligned} \frac{d}{dt} \int V_1 |\mu_t| &\leq -\epsilon \int V_1 |\mu_t| + c_1 \int_{V_0/V_1 \leq \epsilon/K} V_0 |\mu_t| \\ &\leq -\epsilon \int V_1 |\mu_t| + \epsilon_2 \int V_2 |\mu_t|, \end{aligned}$$

where  $\epsilon_2 \ll \epsilon$  as  $\epsilon \rightarrow 0$ . This implies

$$\int V_1 |\mu_{t_n+T}| \leq e^{-\epsilon T} \int V_1 |\mu_{t_n}| + T \epsilon_2 M_2$$

## Subexponential version: proof II

**Second case.** *Estimate for small  $V$ -moment.* Assume now the contrary: there exists  $t_* \in [t_n, t_n + T]$  such that

$$\int V|\mu_{t_*}| \leq \frac{2K}{c_1} \int |\mu_{t_*}|.$$

We take  $t_*$  to be the first time in  $[t_n, t_n + T]$  that satisfies this, and we set  $t_{n+1} := t_* + T$ . With the same bound as before, using the Doeblin assumption we get

$$\|\mu_{t_{n+1}}\|_\beta \leq e^{-\epsilon T} \|\mu_{t_*}\|_\beta + \beta T \epsilon_2 M_2.$$

Since  $\epsilon_2 \ll \epsilon$  as  $\epsilon \rightarrow 0$

One finally gets:

$$r(t) \approx (1 + M_2) \inf_{\epsilon > 0} \left\{ (1 - \epsilon T)^{t/t_0} + T \epsilon_2 \right\}.$$

# A model for neuron populations

Proposed in [Pakdaman, Perthame & Salort 2009]:

$$\left\{ \begin{array}{l} \frac{\partial}{\partial t} n(t, s) + \frac{\partial}{\partial s} n(t, s) + p(N(t), s)n(t, s) = 0, \quad t, s \geq 0, \\ N(t) := n(t, s = 0) = \int_0^{+\infty} p(N(t), s)n(t, s)ds, \quad t > 0, \\ n(t = 0, s) = n_0(s), \quad s > 0. \end{array} \right.$$

- $s \equiv$  time elapsed since last discharge
- $n(t, s) \equiv$  density of neurons which fired  $s$  seconds ago.
- $N(t) \equiv$  total activity at time  $t$
- $p(N, s) \equiv$  probability of firing for a neuron that fired  $s$  seconds ago, given the total activity  $N$ .

# The linear case

If  $p$  does not depend on  $N$  the equation becomes linear:

$$\left\{ \begin{array}{l} \frac{\partial}{\partial t} n(t, s) + \frac{\partial}{\partial s} n(t, s) + p(s)n(t, s) = 0, \quad t, s \geq 0, \\ n(t, s = 0) = \int_0^{+\infty} p(s)n(t, s) ds, \quad t > 0, \\ n(t = 0, s) = n_0(s), \quad s > 0. \end{array} \right.$$

This generates a Markov semigroup in a space of measures (it's linear, conserves mass and positivity).

Notice that the entropy method cannot work in a straightforward way (but see works by **[Perthame & Ryzhik]**, **[Laurençot & Perthame]**, **[Pakdaman, Perthame & Salort]**).

One can apply directly Doeblin's theory to it **[Gabriel 2017]**.

How to show the Doeblin condition:

**Step 1.**  $n$  is larger than the solution  $\tilde{n}$  to

$$\partial_t \tilde{n}(t, s) + \partial_s \tilde{n}(t, s) + p(s) \tilde{n}(t, s) = 0.$$

**Step 2.**  $n$  is larger than the solution to

$$\partial_t n(t, s) + \partial_s n(t, s) + p(s) n(t, s) = 0,$$

$$n(t, 0) = \int_0^{+\infty} p(s) \tilde{n}(t, s) ds.$$



# The weakly nonlinear case

Main assumption:  $|\partial_N p(N, s)|$  small.

Rewrite the nonlinear case as

$$\begin{aligned}\partial_t n &= -\partial_s n - p(N, s)n + N\delta_0 \\ &= L_{N_*} n + (p(N_*, s) - p(N, s))n + \delta_0 \int_0^\infty (p(N, s) - p(N_*, s))n ds \\ &= L_{N_*} n + h\end{aligned}$$

For a given equilibrium  $n_* = n_*(s)$ , call  $N_*$  the corresponding global activity, and  $L_{N_*}$  the operator

$$L_{N_*} n := -\partial_s n(t, s) - p(N_*, s)n(t, s) + \delta_0 \int_0^\infty p(N_*, s)n ds$$

Then:

$$\partial_t(n - n_*) = L_{N_*}(n - n_*) + h$$

So

$$\partial_t(n - n_*) = L_{N_*}(n - n_*) + h,$$

or in other words,

$$n - n_* = S_t(n_0 - n_*) + \int_0^t S_{t-\tau} h(\tau) d\tau$$

This allows us to get an exponential convergence close to  $n_*$ , since

$$\int h = 0, \quad \|h\| \lesssim L \|n - n_*\|$$

# A model for neuron populations with “fatigue”

Proposed in [Pakdaman, Perthame & Salort 2014]:

$$\left\{ \begin{array}{l} \frac{\partial}{\partial t} n(t, s) + \frac{\partial}{\partial s} n(t, s) + p(N(t), s)n(t, s) \\ \quad = \int_0^{+\infty} \kappa(s, u) p(N(t), u) n(t, u) du, \quad u, s, t \geq 0, \\ n(t, s = 0) = 0, \quad N(t) = \int_0^{+\infty} p(N(t), s) n(t, s) ds, \\ n(t = 0, s) = n_0(s) \geq 0, \quad \int_0^{+\infty} n_0(s) ds = 1. \end{array} \right.$$

- $s \equiv$  “state”
- $\kappa(s, u) \equiv$  probability distribution for the landing states when a neuron in state  $u$  jumps to a state  $s$ .

# The linear case

$$\partial_t n(t, s) + \partial_s n(t, s) + p(N(t), s)n(t, s) = \int_0^{+\infty} \kappa(s, u)p(u)n(t, u)du.$$

How to show the Doeblin condition:

**Step 1.**  $n$  is larger than the solution  $\tilde{n}$  to

$$\partial_t \tilde{n}(t, s) + \partial_s \tilde{n}(t, s) + p(s)\tilde{n}(t, s) = 0.$$

**Step 2.**  $n$  is larger than the solution to

$$\partial_t n(t, s) + \partial_s n(t, s) + p(s)n(t, s) = \int_0^{+\infty} \kappa(s, u)p(u)\tilde{n}(t, u)du.$$

We need a condition like:

$$\exists \epsilon, \delta > 0 : \quad \kappa(s, u) \geq \epsilon \mathbf{1}_{[0, \delta]}, \quad \text{for all } s \geq s_*.$$

Thanks for listening!